




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|---|-----------|---|
| (51) International Patent Classification ⁷ : C07D 487/04, A61K 31/53, A61P 25/00, 9/00 | A2 | (11) International Publication Number: WO 00/59907 (43) International Publication Date: 12 October 2000 (12.10.00) |
| (21) International Application Number: PCT/US00/09109 (22) International Filing Date: 6 April 2000 (06.04.00) (30) Priority Data: 60/128,008 6 April 1999 (06.04.99) US (71) Applicant: DU PONT PHARMACEUTICALS COMPANY [US/US]; Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19807 (US). (72) Inventors: GILLIGAN, Paul, J.; 2629 Pennington Drive, Wilmington, DE 19810 (US). WILDE, Richard, G.; 205 Roseman Court, Newark, DE 19711 (US). (74) Agent: RUBIN, Kenneth, B.; Du Pont Pharmaceuticals Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US). | | (81) Designated States: AU, BR, CA, CN, CZ, EE, HU, IL, IN, JP, KR, LT, LV, MX, NO, NZ, PL, RO, SG, SI, SK, TR, UA, VN, ZA, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i> |

(54) Title: PYRAZOLOTRIAZINES AS CRF ANTAGONISTS

(I)

(57) Abstract

The present invention relates to pyrazolotriazines according to formula (I), and stereoisomers, isomers and salts thereof wherein R¹-R⁵ are selected from certain alkyl, aryl and heteroaryl species as defined in the specification wherein all of the compounds are useful as CRF antagonists and are thus useful in the treatment of neurological disorders as well as a multitude of other CRF associated diseases or conditions.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | ML | Mali | TR | Turkey |
| BG | Bulgaria | HU | Hungary | MN | Mongolia | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MR | Mauritania | UA | Ukraine |
| BR | Brazil | IL | Israel | MW | Malawi | UG | Uganda |
| BY | Belarus | IS | Iceland | MX | Mexico | US | United States of America |
| CA | Canada | IT | Italy | NE | Niger | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NL | Netherlands | VN | Viet Nam |
| CG | Congo | KE | Kenya | NO | Norway | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NZ | New Zealand | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | PL | Poland | | |
| CM | Cameroon | KR | Republic of Korea | PT | Portugal | | |
| CN | China | KZ | Kazakhstan | RO | Romania | | |
| CU | Cuba | LC | Saint Lucia | RU | Russian Federation | | |
| CZ | Czech Republic | LI | Liechtenstein | SD | Sudan | | |
| DE | Germany | LK | Sri Lanka | SE | Sweden | | |
| DK | Denmark | LR | Liberia | SG | Singapore | | |
| EE | Estonia | | | | | | |

TITLE

PYRAZOLOTRIAZINES AS CRF ANTAGONISTS

5

FIELD OF THE INVENTION

The present invention relates to novel compounds, compositions, and methods for the treatment of psychiatric disorders and neurological diseases, including major depression, anxiety-related disorders, post-traumatic stress disorder, supranuclear palsy and feeding disorders, as well as treatment of immunological, cardiovascular or heart-related diseases and colonic hypersensitivity associated with psychopathological disturbance and stress. In particular, the present invention relates to novel pyrazolotriazines, pharmaceutical compositions containing such compounds and their use in treating psychiatric disorders, neurological diseases, immunological, cardiovascular or heart-related diseases and colonic hypersensitivity associated with psychopathological disturbance and stress.

BACKGROUND OF THE INVENTION

Corticotropin releasing factor (herein referred to as CRF), a 41 amino acid peptide, is the primary physiological regulator of proopiomelanocortin (POMC) - derived peptide secretion from the anterior pituitary gland [J. Rivier et al., *Proc. Nat. Acad. Sci. (USA)* 80:4851 (1983); W. Vale et al., *Science* 213:1394 (1981)]. In addition to its endocrine role at the pituitary gland, immunohistochemical localization of CRF has demonstrated that the hormone has a broad extrahypothalamic distribution in the central nervous system and produces a wide spectrum of autonomic, electrophysiological and behavioral effects consistent with a neurotransmitter or neuromodulator role in brain [W. Vale et al., *Rec. Prog. Horm. Res.* 39:245 (1983); G.F. Koob, *Persp. Behav. Med.* 2:39 (1985); E.B. De

Souza et al., *J. Neurosci.* 5:3189 (1985)]. There is also evidence that CRF plays a significant role in integrating the response of the immune system to physiological, psychological, and immunological stressors [J.E. Blalock, *Physiological Reviews* 69:1 (1989); J.E. Morley, *Life Sci.* 41:527 (1987)].

Clinical data provide evidence that CRF has a role in psychiatric disorders and neurological diseases including depression, anxiety-related disorders and feeding disorders. A role for CRF has also been postulated in the etiology and pathophysiology of Alzheimer's disease, Parkinson's disease, Huntington's disease, progressive supranuclear palsy and amyotrophic lateral sclerosis as they relate to the dysfunction of CRF neurons in the central nervous system [for review see E.B. De Souza, *Hosp. Practice* 23:59 (1988)].

In affective disorder, or major depression, the concentration of CRF is significantly increased in the cerebral spinal fluid (CSF) of drug-free individuals [C.B. Nemeroff et al., *Science* 226:1342 (1984); C.M. Banki et al., *Am. J. Psychiatry* 144:873 (1987); R.D. France et al., *Biol. Psychiatry* 28:86 (1988); M. Arato et al., *Biol Psychiatry* 25:355 (1989)]. Furthermore, the density of CRF receptors is significantly decreased in the frontal cortex of suicide victims, consistent with a hypersecretion of CRF [C.B. Nemeroff et al., *Arch. Gen. Psychiatry* 45:577 (1988)]. In addition, there is a blunted adrenocorticotropin (ACTH) response to CRF (i.v. administered) observed in depressed patients [P.W. Gold et al., *Am J. Psychiatry* 141:619 (1984); F. Holsboer et al., *Psychoneuroendocrinology* 9:147 (1984); P.W. Gold et al., *New Eng. J. Med.* 314:1129 (1986)]. Preclinical studies in rats and non-human primates provide additional support for the hypothesis that hypersecretion of CRF may be involved in the symptoms seen in human depression [R.M. Sapolsky, *Arch. Gen. Psychiatry* 46:1047 (1989)]. There is preliminary evidence that tricyclic antidepressants

can alter CRF levels and thus modulate the numbers of CRF receptors in brain [Grigoriadis et al., *Neuropsychopharmacology* 2:53 (1989)].

It has also been postulated that CRF has a role in the etiology of anxiety-related disorders. CRF produces anxiogenic effects in animals and interactions between benzodiazepine / non-benzodiazepine anxiolytics and CRF have been demonstrated in a variety of behavioral anxiety models [D.R. Britton et al., *Life Sci.* 31:363 (1982); C.W. Berridge and A.J. Dunn *Regul. Peptides* 16:83 (1986)]. Preliminary studies using the putative CRF receptor antagonist α -helical ovine CRF (9-41) in a variety of behavioral paradigms demonstrate that the antagonist produces "anxiolytic-like" effects that are qualitatively similar to the benzodiazepines [C.W. Berridge and A.J. Dunn *Horm. Behav.* 21:393 (1987), *Brain Research Reviews* 15:71 (1990)].


Neurochemical, endocrine and receptor binding studies have all demonstrated interactions between CRF and benzodiazepine anxiolytics, providing further evidence for the involvement of CRF in these disorders. Chlordiazepoxide attenuates the "anxiogenic" effects of CRF in both the conflict test [K.T. Britton et al., *Psychopharmacology* 86:170 (1985); K.T. Britton et al., *Psychopharmacology* 94:306 (1988)] and in the acoustic startle test [N.R. Swerdlow et al., *Psychopharmacology* 88:147 (1986)] in rats. The benzodiazepine receptor antagonist (Ro15-1788), which was without behavioral activity alone in the operant conflict test, reversed the effects of CRF in a dose-dependent manner while the benzodiazepine inverse agonist (FG7142) enhanced the actions of CRF [K.T. Britton et al., *Psychopharmacology* 94:306 (1988)].

It has been further postulated that CRF has a role in immunological, cardiovascular or heart-related diseases such as hypertension, tachycardia and congestive heart failure, stroke, osteoporosis, premature birth, psychosocial dwarfism, stress-induced


fever, ulcer, diarrhea, post-operative ileus and colonic hypersensitivity associated with psychopathological disturbance and stress.

The mechanisms and sites of action through which the standard anxiolytics and antidepressants produce their therapeutic effects remain to be elucidated. It has been hypothesized however, that they are involved in the suppression of the CRF hypersecretion that is observed in these disorders. Of particular interest is that preliminary studies examining the effects of a CRF receptor antagonist (α -helical CRF9-41) in a variety of behavioral paradigms have demonstrated that the CRF antagonist produces "anxiolytic-like" effects qualitatively similar to the benzodiazepines [for review see G.F. Koob and K.T. Britton, In: *Corticotropin-Releasing Factor: Basic and Clinical Studies of a Neuropeptide*, E.B. De Souza and C.B. Nemeroff eds., CRC Press p221 (1990)].

DuPont Merck PCT application US94/11050 describes corticotropin releasing factor antagonist compounds of the formula:




and their use to treat psychiatric disorders and neurological diseases. Included in the description are fused pyridines and pyrimidines of the formula:




where: V is CR^{1a} or N; Z is CR² or N; A is CR³O or N; and
D is CR²⁸ or N.

WO 98/03510, published in January, 1998, also
describes a series of CRF antagonist compounds having the
5 formula:




wherein z is N or CR² and A is N or CR.

WO 97/29109, published in August, 1997, similarly
10 describes certain CRF antagonist compounds having the
formula:




wherein Ar is phenyl, pyridyl and substituted versions
15 thereof.

WO 98/08847, published March 5, 1998, discloses CRF
antagonist compounds of the formula:




wherein B is selected from a variety of non-aryl groups and R⁵ is selected from certain groups such as phenyl or pyridyl or substituted versions thereof.

WO 99/01454, published on January 14, 1999,
5 discloses CRF antagonist compounds of the formula:




wherein D is an aryl or heteroaryl group and R¹ is selected from certain non-aryl or non-heteroaryl groups.

10 EP 0 269 859 (Ostuka, 1988) discloses pyrazolotriazine compounds of the formula



15 where R¹ is OH or alkanoyl, R² is H, OH, or SH, and R³ is an unsaturated heterocyclic group, naphthyl or substituted phenyl, and states that the compounds have xanthine oxidase inhibitory activity and are useful for treatment of gout.


20 EP 0 594 149 (Ostuka, 1994) discloses pyrazolotriazine and pyrazolopyrimidine compounds of the formula



where A is CH or N, R^0 and R^3 are H or alkyl, and R^1 and R^2 are H, alkyl, alkoxy, alkylthio, nitro, etc., and
 5 states that the compounds inhibit androgen and are useful in treatment of benign prostatic hypertrophy and prostatic carcinoma.


US 3,910,907 (ICN Pharma, 1975) discloses pyrazolotriazines of the formula:

10




where R^1 is CH_3 , C_2H_5 or C_6H_5 , X is H, C_6H_5 , $m-CH_3C_6H_4$, CN, COOEt, Cl, I or Br, Y is H, C_6H_5 , $o-CH_3C_6H_4$, or $p-CH_3C_6H_4$,
 15 and Z is OH, H, CH_3 , C_2H_5 , C_6H_5 , $n-C_3H_7$, $i-C_3H_7$, SH, SCH₃, NHC₄H₉, or $N(C_2H_5)_2$, and states that the compounds are c-AMP phosphodiesterase inhibitors useful as bronchodilators.

US 3,995,039 discloses pyrazolotriazines of the
 20 formula:



where R¹ is H or alkyl, R² is H or alkyl, R³ is H, alkyl, alkanoyl, carbamoyl, or lower alkylcarbamoyl, and R is
 5 pyridyl, pyrimidinyl, or pyrazinyl, and states that the compounds are useful as bronchodilators.


US 5,137,887 discloses pyrazolotriazines of the formula



10

where R is lower alkoxy, and teaches that the compounds are xanthine oxidase inhibitors and are useful for treatment of gout.


15 US 4,892,576 discloses pyrazolotriazines of the formula




20 where X is O or S, Ar is a phenyl, naphthyl, pyridyl or thienyl group, R₆-R₈ are H, alkyl, etc., and R₉ is H,

alkyl, phenyl, etc. The patent states that the compounds are useful as herbicides and plant growth regulants.

US 5,484,760 and WO 92/10098 discloses herbicidal compositions containing, among other things, a herbicidal compound of the formula




where A can be N, B can be CR₃, R₃ can be phenyl or substituted phenyl, etc., R is -N(R₄)SO₂R₅ or -SO₂N(R₆)R₇ and R₁ and R₂ can be taken together to form




where X, Y and Z are H, alkyl, acyl, etc. and D is O or S.

US 3,910,907 and Senga et al., J. Med. Chem., 1982, 25, 243-249, disclose triazolotriazines cAMP phosphodiesterase inhibitors of the formula



where Z is H, OH, CH₃, C₂H₅, C₆H₅, n-C₃H₇, iso-C₃H₇, SH, SCH₃, NH(n-C₄H₉), or N(C₂H₅)₂, R is H or CH₃, and R₁ is CH₃ or C₂H₅. The reference lists eight therapeutic areas where inhibitors of cAMP phosphodiesterase could have utility: asthma, diabetes mellitus, female fertility control, male infertility, psoriasis, thrombosis, anxiety, and hypertension.


WO95/35298 (Otsuka, 1995) discloses pyrazolopyrimidines and states that they are useful as analgesics. The compounds are represented by the formula



5


where Q is carbonyl or sulfonyl, n is 0 or 1, A is a single bond, alkylene or alkenylene, R¹ is H, alkyl, etc., R² is naphthyl, cycloalkyl, heteroaryl, substituted phenyl or phenoxy, R³ is H, alkyl or phenyl, R⁴ is H, alkyl, alkoxycarbonyl, phenylalkyl, optionally phenylthio-substituted phenyl, or halogen, R⁵ and R⁶ are H or alkyl.

EP 0 591 528 (Otsuka, 1991) discloses anti-inflammatory use of pyrazolopyrimidines represented by the formula



where R₁, R₂, R₃ and R₄ are H, carboxyl, alkoxycarbonyl, optionally substituted alkyl, cycloalkyl, or phenyl, R₅ is SR₆ or NR₇R₈, R₆ is pyridyl or optionally substituted phenyl, and R₇ and R₈ are H or optionally substituted phenyl.


Springer et al, J. Med. Chem., 1976, vol. 19, no. 2, 291-296 and Springer U.S. patents 4021,556 and 3,920,652 disclose pyrazolopyrimidines of the formula



5

where R can be phenyl, substituted phenyl or pyridyl, and their use to treat gout, based on their ability to inhibit xanthine oxidase.


10 Joshi et al., J. Prakt. Chemie, 321, 2, 1979, 341-344, discloses compounds of the formula




15 where R¹ is CF₃, C₂F₅, or C₆H₄F, and R² is CH₃, C₂H₅, CF₃, or C₆H₄F.

Maquestiau et al., Bull. Soc. Belg., vol.101, no. 2, 1992, pages 131-136 discloses a pyrazolo[1,5-a]pyrimidine of the formula

20




Ibrahim et al., Arch. Pharm. (weinheim) 320, 487-491 (1987) discloses pyrazolo[1,5-a]pyrimidines of the formula



where R is NH₂ or OH and Ar is 4-phenyl-3-cyano-2-aminopyrid-2-yl.

J. Med. Chem (1982), 25(3), 243-9 discloses compounds of the formula:



wherein R² is H, Ph, Pr, Sme, NH₂Et, NHBu, Net₂, piperidino, OH, NHPr, SH, OCHMe₂, Me, Set, Ome or Opr and R⁴ is H, Br, C₆H₄Me-3, Ph, CN, CO₂Et or Cl.

Other references which disclose azolopyrimidines included EP 0 511 528 (Otsuka, 1992), US 4,997,940 (Dow, 1991), EP 0 374 448 (Nissan, 1990), US 4,621,556 (ICN, 1997), EP 0 531 901 (Fujisawa, 1993), US 4,567,263 (BASF, 1986), EP 0 662 477 (Isagro, 1995), DE 4 243 279 (Bayer, 1994), US 5,397,774 (Upjohn, 1995), EP 0 521 622 (Upjohn, 1993), WO 94/109017 (Upjohn, 1994), J. Med. Chem., 24, 610-613 (1981), and J. Het. Chem., 22, 601 (1985) or others as additionally described herein.

SUMMARY OF THE INVENTION

In accordance with one aspect, the present invention provides novel compounds which bind to corticotropin releasing factor receptors, thereby
5 altering the anxiogenic effects of CRF secretion. The compounds of the present invention are useful for the treatment of psychiatric disorders and neurological diseases, anxiety-related disorders, post-traumatic stress disorder, supranuclear palsy and feeding
10 disorders as well as treatment of immunological, cardiovascular or heart-related diseases and colonic hypersensitivity associated with psychopathological disturbance and stress in mammals.

15 According to another aspect, the present invention provides novel compounds of formula (I) (described below) which are useful as antagonists of the corticotropin releasing factor and which include
20 pyrazolo[1,5-a][1,3,5]triazines and pyrazolo[1,5-a][1,2,4]triazines. The compounds of the present invention exhibit activity as corticotropin releasing factor antagonists and appear to suppress CRF hypersecretion. The present invention also includes
25 pharmaceutical compositions containing such compounds of formula (I), and methods of using such compounds for the suppression of CRF hypersecretion, and/or for the treatment of anxiogenic disorders.

30 According to yet another aspect, the present invention provides novel compounds, pharmaceutical compositions and methods which may be used in the treatment of affective disorder, anxiety, depression, irritable bowel syndrome, post-traumatic stress disorder, supranuclear palsy, immune suppression,
35 Alzheimer's disease, gastrointestinal disease, anorexia nervosa or other feeding disorder, drug or alcohol withdrawal symptoms, drug addiction, inflammatory disorder, fertility problems, disorders, the treatment

of which can be effected or facilitated by antagonizing CRF, including but not limited to disorders induced or facilitated by CRF, or a disorder selected from inflammatory disorders such as rheumatoid arthritis and osteoarthritis, pain, asthma, psoriasis and allergies; generalized anxiety disorder; panic, phobias, obsessive-compulsive disorder; post-traumatic stress disorder; sleep disorders induced by stress; pain perception such as fibromyalgia; mood disorders such as depression, including major depression, single episode depression, recurrent depression, child abuse induced depression, and postpartum depression; dysthemia; bipolar disorders; cyclothymia; fatigue syndrome; stress-induced headache; cancer, human immunodeficiency virus (HIV) infections; neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease and Huntington's disease; gastrointestinal diseases such as ulcers, irritable bowel syndrome, Crohn's disease, spastic colon, diarrhea, and post operative ileus and colonic hypersensitivity associated by psychopathological disturbances or stress; eating disorders such as anorexia and bulimia nervosa; hemorrhagic stress; stress-induced psychotic episodes; euthyroid sick syndrome; syndrome of inappropriate antidiarrhetic hormone (ADH); obesity; infertility; head traumas; spinal cord trauma; ischemic neuronal damage (e.g., cerebral ischemia such as cerebral hippocampal ischemia); excitotoxic neuronal damage; epilepsy; cardiovascular and hear related disorders including hypertension, tachycardia and congestive heart failure; stroke; immune dysfunctions including stress induced immune dysfunctions (e.g., stress induced fevers, porcine stress syndrome, bovine shipping fever, equine paroxysmal fibrillation, and dysfunctions induced by confinement in chickens, sheering stress in sheep or human-animal interaction related stress in dogs); muscular spasms; urinary incontinence; senile dementia of the Alzheimer's type;


multiinfarct dementia; amyotrophic lateral sclerosis;
chemical dependencies and addictions (e.g.,
dependencies on alcohol, cocaine, heroin,
benzodiazepines, or other drugs); drug and alcohol
5 withdrawal symptoms; osteoporosis; psychosocial
dwarfism and hypoglycemia in mammals. The preferred
uses include treatment of depression and anxiety.

The invention further includes use of a compound
of formula I with the variables as recited herein in
10 therapy or the use of a compound of formula I in the
manufacture of a medicament for the treatment of CRF
related diseases or disorders, including anxiety and
depression.

15 According to a still further aspect of the
invention, the compounds provided by this invention
(and especially labelled compounds of this invention)
are also useful as standards and reagents in
determining the ability of a potential pharmaceutical
20 to bind to the CRF receptor.

DETAILED DESCRIPTION OF INVENTION

25 [1] Thus, in a first embodiment, the present invention
provides a novel compound of formula I:



30

or a stereoisomer or pharmaceutically acceptable salt
thereof, wherein:

A equals N or CR⁵;

B equals N or CR⁴ ;

5

provided that both A and B cannot be N or
provided that A can not be CR⁵ and B can not be CR⁴ to
form a pyrazolopyrimidine; and wherein,

10 R¹ is independently selected from the group consisting of

H,

halogen,

CN,

15 C₁₋₆ alkyl,

C₂₋₁₀ alkenyl,

C₂₋₁₀ alkynyl,

C₃₋₆ cycloalkyl,

C₁₋₆ alkyloxy,

20 C₁₋₆ alkylS(O)_n,

-NR^{1a}R^{1b} wherein R^{1a} and R^{1b} are independently selected from

H, C₁₋₄ alkyl, C₃₋₈ cycloalkyl, -C(O)C₁₋₄alkyl,

C₁₋₆ alkylNR^{1a}R^{1b},

NR^{1a}COR^{1b},

25 -C(O)NR^{1a}R^{1b},

-O-C(O)C₁₋₄alkyl,

-XR^{1c} wherein R^{1c} is selected from H or -C₁₋₄ alkylaryl;

X is selected from O or S(O)_n,

30

wherein R¹ is substituted with 0-6 substituents selected
from halogen, C₁₋₄ alkyl, C₃₋₈ cycloalkyl, C₁₋₆ alkyloxy, C₁₋₄
haloalkyl, C₁₋₄alkylamino, C₂₋₈dialkylamino, C₁₋₄alkyloxy, C₁₋₄
alkylthio, C₁₋₄ alkylsulfinyl or C₁₋₄ alkylsulfonyl;

35

R² is selected from the group consisting of

H, OR⁷, SH, NR⁶R⁷, C(OH)R⁶R^{6a}, C(OR⁷)R⁶R^{6a}, S(O)_nR¹³, COR⁷, CO₂R⁷, CHR⁶(OR⁷)R^{6a}, OC(O)R¹³, NO, NO₂, NR⁶C(O)R⁷, N(COR⁷)₂, NR⁶CONR⁶R⁷ or NR⁶CO₂R⁷; or R² is selected from:

- 5 C₁₋₁₀ alkyl,
C₂₋₁₀ alkenyl,
C₂₋₁₀ alkynyl,
C₃₋₈ cycloalkyl,
C₃₋₆ cycloalkyl C₁₋₆ alkyl,
 - 10 C₁₋₁₀ alkyloxy,
C₁₋₁₀ alkyloxyC₁₋₁₀ alkyl,
-SO₂-C₁₋₁₀alkyl
-SO₂R^{2a} wherein R^{2a} is aryl,
-SO₂R^{2b} wherein R^{2b} is heteroaryl,
 - 15 -NR^{2c}R^{2d} wherein R^{2c} and R^{2d} are independently selected from
H, C₁₋₈ alkyl, S(O)_nC₁₋₄alkyl, C(O)NR^{2c}R^{2d}, CO₂C₁₋₄alkyl,
C₃₋₈ cycloalkyl, C₁₋₆ alkyloxyC₁₋₆ alkyl, -C(O)C₁₋₄alkyl
or R^{2c} and R^{2d} may join to form a heterocyclic ring
having 0-3 heteroatoms selected from O, N or S,
 - 20 - halogen,
-CN,
-C(O)-L wherein L is selected from H, NR^{2c}R^{2d}, C₁₋₆ alkyl or
OC₁₋₄ alkyl, O(CH₂)_nOR wherein R is C₁₋₃ alkyl, O(CH₂)_n-
25 NR^{2c}R^{2d}, OH, C(O)OC₁₋₆alkyl, or aryl or heteroaryl
wherein m is 1-4; or

-OC(O)-M wherein M is selected from C₁₋₄ alkyl, C₁₋₄
haloalkyl, C₂₋₈ alkoxyalkyl, C₃₋₈cycloalkyl, C₄₋₁₂
30 cycloalkylalkyl, aryl, C₁₋₆ alkylaryl, heteroaryl, C₁₋₆
alkylheteroaryl;
- n is 0, 1 or 2; and wherein
- 35 R² is substituted with 0-3 substituents independently
selected from R', R'', R''' wherein R', R'' and R''' are
independently selected from C₁₋₆ alkyl, C₃₋₇ cycloalkyl,

hydroxyC₁₋₆ alkyl, C₁₋₆ alkyloxyC₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ alkyloxy, hydroxy, or

R² is substituted with 0-3 substituents independently
5 selected from:

halogen,

-CN,

-S(O)_nR^{2e} wherein R^{2e} is selected from C₁₋₄ alkyl, C₁₋₄
10 haloalkyl, C₁₋₄ alkyloxy C₁₋₄ alkyl, C₃₋₆ cycloalkyl;

-COR^{2f} wherein R^{2f} is selected from H, C₁₋₄ alkyl, C₁₋₄
haloalkyl, C₁₋₄ alkyloxy C₁₋₄ alkyl, C₃₋₆
cycloalkyl, and C₃₋₆ cycloalkylC₁₋₄ alkyl;

15

-CO₂R^{2f},

-NR^{2g}COR^{2f} wherein R^{2g} is selected from H, C₁₋₆ alkyl, C₃₋₇
cycloalkyl, C₃₋₆ cycloalkyl C₁₋₆ alkyl;

-N(COR^{2f})₂,

20 -NR^{2g}CONR^{2f}R^{2h}, wherein R^{2h} is selected from H, C₁₋₆ alkyl,
C₁₋₄ haloalkyl, C₁₋₄ alkoxy C₁₋₄ alkyl,
C₃₋₆ cycloalkyl and C₃₋₆ cycloalkylC₁₋₆
alkyl;

25 -NR^{2g}CO₂R^{2o},

-CONR^{2g}R^{2h},

1-morpholinyl,

1-piperidinyl,

1-piperazinyl,

30 and

C₃₋₈ cycloalkyl wherein 0-1 carbon atoms in the C₄₋₈
cycloalkyl is replaced by a group selected from

-O-, -S(O)_n-, -NR^{2g}-, -NCO₂R^{2o}, -NCOR^{2o},

and -NSO₂R^{2o}; and wherein N₄ in

35 1-piperazinyl is substituted with 0-1
substituents selected from R^{2g}, CO₂R^{2o}, COR^{2e} and
SO₂R^{2o}; or

the group R^{2i} , R^{2j} , R^{2k} , C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, Br, Cl, F, I, C_{1-4} haloalkyl, $-OR^{2g}$, $-NR^{2g}R^{2h}$, $-C_{1-6}$ alkyl- OR^{2g} , and C_{3-8} cycloalkyl which is substituted with 0-1 R^{2i} and in which 0-1 carbons of C_{4-8} cycloalkyl is replaced by -O-, wherein

R^{2i} is selected from aryl wherein aryl includes phenyl, naphthyl, indanyl and indenyl, each R^{2i} being substituted with 0-1 OR^{2m} and 0-5 substituents independently selected from the group C_{1-6} alkyl, C_{3-6} cycloalkyl, Br, Cl, F, I, C_{1-4} haloalkyl, $-CN$, nitro, $-SH$, $-S(O)_nR^{2n}$, $-COR^{2m}$, $-OC(O)R^{2n}$, $-NR^{2g}COR^{2m}$, $-N(COR^{2m})_2$, $-NR^{2g}CONR^{2o}R^{2p}$, $-NR^{2g}CO_2R^{2n}$, $-NR^{2o}R^{2p}$ and $-CONR^{2o}R^{2p}$;

R^{2j} is selected from heteroaryl wherein heteroaryl includes pyridyl, pyrimidinyl, triazinyl, furanyl, quinolinyl, isoquinolinyl, thienyl, imidazolyl, thiazolyl, indolyl, pyrrolyl, oxazolyl, benzofuranyl, benzothienyl, benzothiazolyl, benzoxazolyl, isoxazolyl, pyrazolyl, triazolyl, tetrazolyl, indazolyl, 2,3-dihydrobenzofuranyl, 2,3-dihydrobenzothienyl, 2,3-dihydrobenzothienyl-s-oxide, 2,3-dihydro-benzothienyl-S-dioxide, indolinyl, benzoxazolin-2-onyl, benzodioxolanyl and benzodioxane, each heteroaryl being substituted on 0-4 carbon atoms with a substituent independently selected from the group C_{1-6} alkyl, C_{3-6} cycloalkyl, Br, Cl, F, I, C_{1-4} haloalkyl, $-CN$, nitro, OR^{2m} , $-SH$, $-S(O)_nR^{2h}$, $-COR^{2m}$, $-OC(O)R^{2h}$, $-NR^{2g}COR^{2m}$, $-N(COR^{2m})_2$, $-NR^{2g}CONR^{2o}R^{2p}$, $-NR^{2g}CO_2R^{2h}$, $-NR^{2o}R^{2p}$ and $-CONR^{2o}R^{2p}$ and each heteroaryl being substituted on any nitrogen atom with 0-1 substituents selected from the group R^{2g} , CO_2R^{2e} , COR^{2e} and SO_2R^{2e} ;

R^{2k} is heterocyclyl which is a saturated or partially saturated heteroaryl as defined for R^{2j} , each heterocyclyl being substituted on 0-4 carbon atoms with a substituent independently selected from the group C_{1-6} alkyl, C_{3-6} cycloalkyl, Br, Cl, F, I, C_{1-4} haloalkyl, $-CN$, nitro, $-OR^{2m}$,

-SH, $-S(O)_n R^{2h}$, $-COR^{2m}$, $-OC(O)R^{2h}$, $-NR^{2g}COR^{2m}$, $-N(COR^{2m})_2$,
 $-NR^{2g}CONR^{2o}R^{2p}$, $NR^{2g}CO_2R^{2h}$, $-NR^{2o}R^{2p}$ and $-CONR^{2o}R^{2p}$ and each
heterocyclyl being substituted on any nitrogen atom with
0-1 substituents selected from the group R^{2f} , CO_2R^{2e} , COR^{2e}
5 and SO_2R^{2e} ;

wherein

R^{21} is H, C_{1-4} alkyl, C_{3-6} cycloalkyl- C_{1-4} alkyl and C_{3-8}
10 cycloalkyl;

R^{2m} is H, C_{1-6} alkyl, C_{3-6} cycloalkyl C_{1-6} alkyl, C_{1-2}
alkyloxy C_{1-2} alkyl, C_{1-4} haloalkyl, $R^{2q}S(O)_n-C_{1-4}$ alkyl
and $R^{2r}R^{2s}N-C_{2-4}$ alkyl;
15

R^{2n} is H, C_{1-6} alkyl, C_{3-10} cycloalkyl, C_{3-6} cycloalkyl-
 C_{1-6} alkyl, C_{1-2} alkyloxy C_{1-2} alkyl, and C_{1-4} haloalkyl;

R^{2o} and R^{2p} are independently selected at each occurrence
20 from H, C_{1-6} alkyl, C_{3-10} cycloalkyl, C_{3-6} cycloalkyl C_{1-6} alkyl
and C_{1-4} haloalkyl;

R^{2q} is selected from C_{1-6} alkyl, C_{1-4} haloalkyl, C_{1-4} alkoxy-
 C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkyl- C_{1-6} alkyl, aryl,
25 aryl(C_{1-4} alkyl), heteroaryl and heteroaryl (C_{1-4} alkyl)-
and benzyl, each benzyl being substituted on the aryl
moiety with 0-1 substituents selected from the group C_{1-4}
alkyl, Br, Cl, F, I, C_{1-4} haloalkyl, nitro, C_{1-4} alkoxy C_{1-4}
haloalkoxy, and dimethylamino;

30 $R^{2r}R^{2s}$ taken together with the N form 1-pyrrolidinyl, 1-
morpholinyl, 1-piperidinyl or 1-piperazinyl wherein N_4 in
1-piperiazinyl is substituted with 0-1 substituents
selected from the group R^{2t} , CO_2R^{2q} , COR^{2q} and SO_2R^{2q} ;

35 R^{2t} is selected from H, C_{1-4} alkyl, C_{1-4} haloalkyl, C_{1-4} alkoxy
 $-C_{1-4}$ alkyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkyl - C_{1-6} alkyl,

aryl, aryl (C_{1-4} alkyl)-, heteroaryl and heteroaryl (C_{1-4} alkyl);

R^3 is selected from an aryl or heteroaryl group attached
5 through an unsaturated carbon atom;

aryl is selected from phenyl, naphthyl, indanyl and indenyl, each aryl being substituted with 0-5 substituents independently selected at each occurrence
10 from C_{1-6} alkyl, C_{3-6} cycloalkyl, methylenedioxy, C_{1-4} alkyloxy- C_{1-4} alkyloxy, $-OR^{2m}$, Br, Cl, F, I, C_{1-4} haloalkyl, $-CN$, $-NO_2$, $-SH$, $-S(O)_nR^{2n}$, $-COR^{2m}$, $-CO_2R^{2m}$, $-OC(O)R^{2n}$, $-NR^{2g}COR^{2m}$, $-N(COR^{2m})_2$, $-NR^{2g}CONR^{2o}R^{2p}$, $-NR^{2g}CO_2R^{2h}$, $-NR^{2o}R^{2p}$ and $CONR^{2o}R^{2p}$;

15 heteroaryl is selected from the group pyridyl, pyrimidyl, triazinyl, furanyl, quinoliny, isoquinoliny, thienyl, imidazolyl, thiazolyl, indolyl, pyrrolyl, oxazolyl, benzofuranyl, benzothienyl, benzothiazolyl, benzoxazolyl,
20 isoxazolyl, triazolyl, tetrazolyl, indazolyl, 2,3-dihydrobenzo-furanyl, 2,3-dihydrobenzothienyl, 2,3-dihydro-benzothienyl-S-oxide, 2,3-dihydrobenzothienyl-s-dioxide, indoliny, benzoxazolin-2-on-yl, benzodioxolanyl and benzodioxane, each heteroaryl being substituted at 0-
25 4 carbon atoms with a substituent independently selected at each occurrence from the group C_{1-6} alkyl, C_{3-6} cycloalkyl, Br, F, I, C_{1-4} haloalkyl, $-CN$, $NR^{2g}R^{2h}$, nitro, $-OR^{2m}$, $-SH$, $-S(O)_nR^{2n}$, COR^{2m} , $-CO_2R^{2m}$, $-OC(O)R^{2n}$, $-NR^{2g}COR^{2m}$, $-N(COR^{2m})_2$, $-NR^{2g}CONR^{2o}R^{2p}$ and each heteroaryl being
30 substituted at any nitrogen atom with 0-1 substituents selected from the group R^{2g} , CO_2R^{3a} , COR^{3a} and SO_2R^{3a} wherein,

R^{3a} is selected from the group C_{1-6} alkyl, C_{1-4} cycloalkyl- C_{1-6} alkyl and benzyl, each benzyl being substituted on the
35 aryl moiety with 0-1 substituents selected from the group C_{1-4} alkyl, Br, Cl, F, I, C_{1-4} haloalkyl, nitro, C_{1-4} alkoxy, C_{1-4} haloalkoxy, and dimethylamino;

- R^4 and R^5 are independently selected at each occurrence from H, Br, Cl, F, I, -CN, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-8} cycloalkyl, C_{1-6} alkyloxy, C_{1-6} alkylthio, C_{1-6} alkylsulfinyl, C_{1-6} alkylsulfonyl, amino, C_{1-4} alkylamino, (C₁₋₄ alkyl)₂ amino and phenyl, each phenyl is substituted with 0-3 groups selected from the group consisting of C_{1-4} alkyl, C_{3-8} cycloalkyl, Br, Cl, F, I, -C(O)H, C_{1-4} haloalkyl, nitro, C_{1-4} alkoxy, C_{1-4} haloalkoxy, C_{1-4} alkylthio, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, C_{1-6} alkylamino and (C₁₋₄ alkyl)₂ amino and wherein R^4 and R^5 non-phenyl groups may be substituted with 0-5 substituents selected from OH, halogen, -C(O)H, -OC₁₋₆-alkyl and C_{1-6} haloalkyl, C_{1-6} alkyl, C_{3-7} c-alkyl, C_{1-6} alkyl(OH)_nCO₂R wherein R is H or C_{1-6} alkyl, C_{1-6} alkyl(OH)_n, wherein n is 0-3 or R^4 and R^5 may join together to form a C_{3-6} alkylene chain;

- R^6 , R^{6a} and R^7 are independently selected from:
H, C_{1-10} alkyl, C_{3-10} cycloalkyl, C_{3-10} alkenyl, C_{3-10} alkynyl, C_{1-10} haloalkyl, C_{2-8} alkoxyalkyl, C_{4-12} cycloalkylalkyl, C_{5-10} cycloalkenyl, C_{6-14} cycloalkenylalkyl;

- R^6 , R^{6a} and R^7 are substituted with 0-6 substituents independently selected from halogen, C_{1-4} alkyl, C_{3-8} cycloalkyl, C_{1-6} alkyloxy, C_{1-4} haloalkyl;


with the proviso that the compounds of Formula I with R^1 , R^2 , R^3 , R^4 and R^5 as specifically defined below are excluded:

- (a) a compound of formula I wherein A = CR⁵ with R⁵ o-hydroxyphenyl, B = N, R³ = o-hydroxyphenyl, R¹=SMe and R²=CN (Registry Reference 23/IS98062) and
- (b) a compound of formula I wherein A=CR⁵, R⁵=CH₃, B = N, R¹ = Ph, R² = Br and R³ is Ph;

- (c) a compound of formula I wherein $A = CR^5$, $R^5 = p\text{-Cl-phenyl}$, $B = N$, $R^1 = \text{Me}$, $R^2 = H$ and $R^3 = p\text{-CF}_3\text{-phenyl}$ (Registry reference 152/IS98062);
- 5 (d) a compound of formula I wherein $A = CR^5$, $R^5 = \text{phenyl}$, $B = N$, $R^1 = \text{Me}$, $R^2 = H$ and $R^3 = p\text{-CF}_3\text{-phenyl}$ (Registry reference 153/IS98062);
- (e) a compound of formula I wherein $A = CR^5$, $R^5 = \text{ethyl}$,
10 $B = N$, $R^1 = \text{Me}$, $R^2 = H$ and $R^3 = \text{N-methyl-piperiazin-N-yl}$ (registry reference 184/IS98062);
- (f) a compound of formula I wherein $A = CR^5$, R^5 is $p\text{-Cl-Ph}$, $R^1 = H$, $R^2 = H$ and $R^3 = p\text{-CF}_3\text{-Ph}$ (Registry reference
15 9/IS98179);
- (g) a compound of formula I wherein $A = CR^5$, $R^5 = p\text{-Cl-Ph}$, $R^1 = \text{CH}_3$, $R^2 = H$, $R^3 = p\text{-CF}_3\text{-Ph}$ (Registry reference 10/IS98179);
- 20 (h) a compound of formula I wherein $A = CR^5$, $R^5 = \text{Ph}$, $R^1 = \text{Me}$, $R^2 = H$, $R^3 = p\text{-CF}_3\text{-Ph}$ (Registry reference 11/IS98179);
- (i) a compound of formula I wherein $A = CR^5$, $R^5 = \text{Ph}$, $R^1 = H$, $R^2 = H$, $R^3 = p\text{-CF}_3\text{-Ph}$ (Registry reference 12/IS98179);
25
- (j) a compound of formula I wherein $A = CR^5$, $R^1 = \text{Ph}$ and R^2 is H , Br , CN , CO_2Et or Cl (J. Med. Chem. (1982), 25(3), 243-9;
- 30 (k) a compound of formula I wherein $A = CR^5$, $R^5 = \text{CH}_3$, C_2H_5 or Ph , $R^1 = H$, $R^2 = H$ and $R^3 = \text{Ph}$ (US 3,910,907).

[1'] The present invention preferably relates to
a novel compound of formula I:

35



or a stereoisomer or pharmaceutically acceptable salt thereof, wherein:

5

A equals N or CR⁵;

B equals N or CR⁴ ;

- 10 provided that both A and B cannot be N or
provided that A can not be CR⁵ and B can not be CR⁴ to
form a pyrazolopyrimidine; and wherein,

R¹ is independently selected from the group consisting of

15

H,

halogen,

CN,

C₁₋₆ alkyl,

20 C₂₋₁₀ alkenyl,

C₂₋₁₀ alkynyl,

C₃₋₆ cycloalkyl,

C₁₋₆ alkyloxy,

C₁₋₆ alkylS(O)_n,

25 -NR^{1a}R^{1b} wherein R^{1a} and R^{1b} are independently selected from

H, C₁₋₄ alkyl, C₃₋₈ cycloalkyl, -C(O)C₁₋₄alkyl,

C₁₋₆ alkylNR^{1a}R^{1b},

NR^{1a}COR^{1b},

-C(O)NR^{1a}R^{1b},

30 -O-C(O)C₁₋₄alkyl,

-XR^{1c} wherein R^{1c} is selected from H or -C₁₋₄ alkylaryl;

X is selected from 0 or $S(O)_n$,

wherein R^1 is substituted with 0-6 substituents selected from halogen, C_{1-4} alkyl, C_{3-8} cycloalkyl, C_{1-6} alkyloxy, C_{1-4} haloalkyl, C_{1-4} alkylamino, C_{2-8} dialkylamino, C_{1-4} alkyloxy, C_{1-4} alkylthio, C_{1-4} alkylsulfinyl or C_{1-4} alkylsulfonyl;

R^2 is selected from the group consisting of OR^7 , SH , NR^6R^7 , $C(OH)R^6R^{6a}$, $C(OR^7)R^6R^{6a}$, $S(O)_nR^{13}$, COR^7 , CO_2R^7 , $CHR^6(OR^7)R^{6a}$, $OC(O)R^{13}$, NO , NO_2 , $NR^6C(O)R^7$, $N(COR^7)_2$, $NR^6CONR^6R^7$ or $NR^6CO_2R^7$; or R^2 is selected from:

- C_{1-10} alkyl,
- C_{2-10} alkenyl,
- 15 C_{2-10} alkynyl,
- C_{3-8} cycloalkyl,
- C_{3-6} cycloalkyl C_{1-6} alkyl,
- C_{1-10} alkyloxy,
- C_{1-10} alkyloxy C_{1-10} alkyl,
- 20 $-SO_2-C_{1-10}$ alkyl
- $-SO_2R^{2a}$ wherein R^{2a} is aryl,
- $-SO_2R^{2b}$ wherein R^{2b} is heteroaryl,
- $-NR^{2c}R^{2d}$ wherein R^{2c} and R^{2d} are independently selected from
- 25 H , C_{1-8} alkyl, $S(O)_nC_{1-4}$ alkyl, $C(O)NR^{2c}R^{2d}$, CO_2C_{1-4} alkyl,
- C_{3-8} cycloalkyl, C_{1-6} alkyloxy C_{1-6} alkyl, $-C(O)C_{1-4}$ alkyl
- or R^{2c} and R^{2d} may join to form a heterocyclic ring
- having 0-3 heteroatoms selected from O, N or S,
- 30 $-C(O)-L$ wherein L is selected from H , $NR^{2c}R^{2d}$, C_{1-6} alkyl
- $O(CH_2)_mOR$ wherein R is C_{1-3} alkyl, $O(CH_2)_m-NR^{2c}R^{2d}$, OH ,
- $C(O)OC_{1-6}$ alkyl, or aryl or heteroaryl wherein m is 1-4; or
- $-OC(O)-M$ wherein M is selected from C_{1-4} alkyl, C_{1-4}
- 35 haloalkyl, C_{2-8} alkoxyalkyl, C_{3-6} cycloalkyl, C_{4-12}
- cycloalkylalkyl, aryl, C_{1-6} alkylaryl, heteroaryl, C_{1-6}
- alkylheteroaryl;

n is 0, 1 or 2; and wherein

R^2 is substituted with 0-3 substituents independently
selected from R' , R'' , R''' wherein R' , R'' and R''' are
5 independently selected from C_{1-6} alkyl, C_{3-7} cycloalkyl,
hydroxy C_{1-6} alkyl, C_{1-6} alkyloxy C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6}
alkynyl, C_{1-6} alkyloxy, hydroxy, or

R^2 is substituted with 0-3 substituents independently
10 selected from:

halogen,

-CN,

-S(O) $_n$ R^{2o} wherein R^{2o} is selected from C_{1-4} alkyl, C_{1-4}
15 haloalkyl, C_{1-4} alkyloxy C_{1-4} alkyl, C_{3-6} cycloalkyl;

-COR 2f wherein R^{2f} is selected from H, C_{1-4} alkyl, C_{1-4}
haloalkyl, C_{1-4} alkyloxy C_{1-4} alkyl, C_{3-6}
cycloalkyl, and C_{3-6} cycloalkyl C_{1-4} alkyl;

20

-CO $_2$ R^{2f} ,

-NR 2g COR 2f wherein R^{2g} is selected from H, C_{1-6} alkyl, C_{3-7}
cycloalkyl, C_{3-6} cycloalkyl C_{1-6} alkyl;

-N(COR 2f) $_2$,

25 -NR 2g CONR 2f R^{2h} , wherein R^{2h} is selected from H, C_{1-6} alkyl,
 C_{1-4} haloalkyl, C_{1-4} alkoxy C_{1-4} alkyl,
 C_{3-6} cycloalkyl and C_{3-6} cycloalkyl C_{1-6}
alkyl;

30 -NR 2g CO $_2$ R^{2o} ,

-CONR 2g R^{2h} ,

1-morpholinyl,

1-piperidinyl,

1-piperazinyl,

35 and

C_{3-8} cycloalkyl wherein 0-1 carbon atoms in the C_{4-8}
cycloalkyl is replaced by a group selected from
-O-, -S(O) $_n$ -, -NR 2g -, -NCO $_2$ R^{2o} , -NCOR 2o ,

and $-\text{NSO}_2\text{R}^{2e}$; and wherein N_i in
 1-piperazinyl is substituted with 0-1
 substituents selected from R^{2g} , CO_2R^{2e} , COR^{2e} and
 SO_2R^{2e} ; or

5 the group R^{2i} , R^{2j} , R^{2k} , C_{1-6} alkyl, C_{2-8}
 alkenyl, C_{2-8} alkynyl, Br, Cl, F, I, C_{1-4} haloalkyl, $-\text{OR}^{2g}$,
 $-\text{NR}^{2g}\text{R}^{2h}$, $-\text{C}_{1-6}$ alkyl- OR^{2g} , and C_{3-8} cycloalkyl which is
 substituted with 0-1 R^{2i} and in which 0-1 carbons of C_{4-8}
 10 cycloalkyl is replaced by -O-, wherein

R^{2i} is selected from aryl wherein aryl includes
 phenyl, naphthyl, indanyl and indenyl, each
 R^{2i} being substituted with 0-1 OR^{2m} and 0-5

15 substituents independently selected from the group C_{1-6}
 alkyl, C_{3-6} cycloalkyl, Br, Cl, F, I, C_{1-4} haloalkyl, -CN,
 nitro, -SH, $-\text{S}(\text{O})_n\text{R}^{2n}$, $-\text{COR}^{2m}$, $-\text{OC}(\text{O})\text{R}^{2n}$, $-\text{NR}^{2g}\text{COR}^{2m}$, -
 $\text{N}(\text{COR}^{2m})_2$,
 $-\text{NR}^{2g}\text{CONR}^{2o}\text{R}^{2p}$, $-\text{NR}^{2g}\text{CO}_2\text{R}^{2n}$, $-\text{NR}^{2o}\text{R}^{2p}$ and $-\text{CONR}^{2o}\text{R}^{2p}$;

20 R^{2j} is selected from heteroaryl wherein heteroaryl
 includes pyridyl, pyrimidinyl, triazinyl, furanyl,
 quinolinyl, isoquinolinyl, thienyl, imidazolyl,
 thiazolyl, indolyl, pyrrolyl, oxazolyl, benzofuranyl,
 25 benzothienyl, benzothiazolyl, benzoxazolyl, isoxazolyl,
 pyrazolyl, triazolyl, tetrazolyl, indazolyl, 2,3-
 dihydrobenzofuranyl, 2,3-dihydrobenzothienyl, 2,3-
 dihydrobenzothienyl-s-oxide, 2,3-dihydro-benzothienyl-S-
 dioxide, indolinyl, benzoxazolin-2-onyl, benzodioxolanyl
 30 and benzodioxane, each heteroaryl being substituted on 0-
 4 carbon atoms with a substituent independently selected
 from the group C_{1-6} alkyl, C_{3-6} cycloalkyl, Br, Cl, F, I, C_{1-4}
 haloalkyl, -CN, nitro, OR^{2m} , -SH, $-\text{S}(\text{O})_n\text{R}^{2h}$, $-\text{COR}^{2m}$, -
 $\text{OC}(\text{O})\text{R}^{2h}$, $-\text{NR}^{2g}\text{COR}^{2m}$, $-\text{N}(\text{COR}^{2m})_2$, $-\text{NR}^{2g}\text{CONR}^{2o}\text{R}^{2p}$, $-\text{NR}^{2g}\text{CO}_2\text{R}^{2h}$, -
 35 $\text{NR}^{2o}\text{R}^{2p}$ and $-\text{CONR}^{2o}\text{R}^{2p}$ and each heteroaryl being substituted
 on any nitrogen atom with 0-1 substituents selected from
 the group R^{2g} , CO_2R^{2e} , COR^{2e} and SO_2R^{2e} ;

R^{2k} is heterocyclyl which is a saturated or partially saturated heteroaryl as defined for R^{2j} , each heterocyclyl being substituted on 0-4 carbon atoms with a substituent independently selected from the group C_{1-6} alkyl, C_{3-6} cycloalkyl, Br, Cl, F, I, C_{1-4} haloalkyl, -CN, nitro, -OR^{2m}, -SH, -S(O)_nR^{2h}, -COR^{2m}, -OC(O)R^{2h}, -NR^{2g}COR^{2m}, -N(COR^{2m})₂, -NR^{2g}CONR^{2o}R^{2p}, NR^{2g}CO₂R^{2h}, -NR^{2o}R^{2p} and -CONR^{2o}R^{2p} and each heterocyclyl being substituted on any nitrogen atom with 0-1 substituents selected from the group R^{2f}, CO₂R^{2c}, COR^{2e} and SO₂R^{2e};

wherein

R^{21} is H, C_{1-4} alkyl, C_{3-6} cycloalkyl- C_{1-4} alkyl and C_{3-8} cycloalkyl;

R^{2m} is H, C_{1-6} alkyl, C_{3-6} cycloalkyl C_{1-6} alkyl, C_{1-2} alkyloxy C_{1-2} alkyl, C_{1-4} haloalkyl, R^{2a}S(O)_n- C_{1-4} alkyl and R^{2r}R^{2e}N- C_{2-4} alkyl;

R^{2n} is H, C_{1-6} alkyl, C_{3-10} cycloalkyl, C_{3-6} cycloalkyl- C_{1-6} alkyl, C_{1-2} alkyloxy C_{1-2} alkyl, and C_{1-4} haloalkyl;

R^{2o} and R^{2p} are independently selected at each occurrence from H, C_{1-6} alkyl, C_{3-10} cycloalkyl, C_{3-6} cycloalkyl C_{1-6} alkyl and C_{1-4} haloalkyl;

R^{2a} is selected from C_{1-6} alkyl, C_{1-4} haloalkyl, C_{1-4} alkoxy- C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkyl- C_{1-6} alkyl, aryl, aryl(C_{1-4} alkyl), heteroaryl and heteroaryl (C_{1-4} alkyl)- and benzyl, each benzyl being substituted on the aryl moiety with 0-1 substituents selected from the group C_{1-4} alkyl, Br, Cl, F, I, C_{1-4} haloalkyl, nitro, C_{1-4} alkoxy C_{1-4} haloalkoxy, and dimethylamino;

R^{2r}R^{2o} taken together with the N form 1-pyrrolidinyl, 1-morpholinyl, 1-piperidinyl or 1-piperazinyl wherein N₄ in

1-piperiazinyl is substituted with 0-1 substituents selected from the group R^{2t} , CO_2R^{2q} , COR^{2q} and SO_2R^{2q} ;

R^{2t} is selected from H, C_{1-4} alkyl, C_{1-4} haloalkyl, C_{1-4} alkoxy
5 $-C_{1-4}$ alkyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkyl - C_{1-6} alkyl, aryl, aryl (C_{1-4} alkyl)-, heteroaryl and heteroaryl (C_{1-4} alkyl);

R^3 is selected from an aryl or heteroaryl group attached
10 through an unsaturated carbon atom;

aryl is selected from phenyl, naphthyl, indanyl and indenyl, each aryl being substituted with 0-5 substituents independently selected at each occurrence
15 from C_{1-6} alkyl, C_{3-6} cycloalkyl, methylenedioxy, C_{1-4} alkyloxy- C_{1-4} alkyloxy, $-OR^{2m}$, Br, Cl, F, I, C_{1-4} haloalkyl, $-CN$, $-NO_2$, $-SH$, $-S(O)_nR^{2n}$, $-COR^{2m}$, $-CO_2R^{2m}$, $-OC(O)R^{2n}$, $-NR^{2q}COR^{2m}$, $-N(COR^{2m})_2$, $-NR^{2q}CONR^{2o}R^{2p}$, $-NR^{2q}CO_2R^{2h}$, $-NR^{2o}R^{2p}$ and $CONR^{2o}R^{2p}$;

20 heteroaryl is selected from the group pyridyl, pyrimidyl, triazinyl, furanyl, quinolinyl, isoquinolinyl, thienyl, imidazolyl, thiazolyl, indolyl, pyrrolyl, oxazolyl, benzofuranyl, benzothienyl, benzothiazolyl, benzoxazolyl,
25 isoxazolyl, triazolyl, tetrazolyl, indazolyl, 2,3-dihydrobenzo-furanyl, 2,3-dihydrobenzothienyl, 2,3-dihydro-benzothienyl-S-oxide, 2,3-dihydrobenzothienyl-s-dioxide, indolinyl, benzoxazolin-2-on-yl, benzodioxolanyl and benzodioxane, each heteroaryl being substituted at 0-
30 4 carbon atoms with a substituent independently selected at each occurrence from the group C_{1-6} alkyl, C_{3-6} cycloalkyl, Br, F, I, C_{1-4} haloalkyl, $-CN$, $NR^{2q}R^{2h}$, nitro, $-OR^{2m}$, $-SH$, $-S(O)_nR^{2n}$, COR^{2m} , $-CO_2R^{2m}$, $-OC(O)R^{2n}$, $-NR^{2q}COR^{2m}$, $-N(COR^{2m})_2$, $-NR^{2q}CONR^{2o}R^{2p}$ and each heteroaryl being
35 substituted at any nitrogen atom with 0-1 substituents selected from the group R^{2q} , CO_2R^{3a} , COR^{3a} and SO_2R^{3a} wherein,



R^{3a} is selected from the group C₁₋₆ alkyl, C₁₋₄ cycloalkyl-C₁₋₆ alkyl and benzyl, each benzyl being substituted on the aryl moiety with 0-1 substituents selected from the group C₁₋₄ alkyl, Br, Cl, F, I, C₁₋₄ haloalkyl, nitro, C₁₋₄ alkoxy, C₁₋₄ haloalkoxy, and dimethylamino;

R⁴ and R⁵ are independently selected at each occurrence from H, Br, Cl, F, I, -CN, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₈ cycloalkyl, C₁₋₆ alkyloxy, C₁₋₆ alkylthio, C₁₋₆ alkylsulfanyl, C₁₋₆ alkylsulfonyl, amino, C₁₋₄ alkylamino, (C₁₋₄ alkyl)₂ amino and phenyl, each phenyl is substituted with 0-3 groups selected from the group consisting of C₁₋₇ alkyl, C₃₋₈ cycloalkyl, Br, Cl, F, I, -C(O)H, C₁₋₄ haloalkyl, nitro, C₁₋₄ alkoxy, C₁₋₄ haloalkoxy, C₁₋₄ alkylthio, C₁₋₄ alkylsulfanyl, C₁₋₄ alkylsulfonyl, C₁₋₆ alkylamino and (C₁₋₄ alkyl)₂ amino and wherein R⁴ and R⁵ non-phenyl groups may be substituted with 0-5 substituents selected from OH, halogen, -C(O)H, -OC₁₋₆-alkyl and C₁₋₆ haloalkyl, C₁₋₆ alkyl, C₃₋₇ c-alkyl, C₁₋₆ alkyl(OH)_nCO₂R wherein R is H or C₁₋₆ alkyl, C₁₋₆ alkyl(OH)_n, wherein n is 0-3 or R⁴ and R⁵ may join together to form a C₃₋₆ alkylene chain;

R⁶, R^{6a} and R⁷ are independently selected from:
H, C₁₋₁₀ alkyl, C₃₋₁₀ cycloalkyl, C₃₋₁₀ alkenyl, C₃₋₁₀ alkynyl, C₁₋₁₀ haloalkyl, C₂₋₈ alkoxyalkyl, C₄₋₁₂ cycloalkylalkyl, C₅₋₁₀ cycloalkenyl, C₆₋₁₄ cycloalkenylalkyl;

R⁶, R^{6a} and R⁷ are substituted with 0-6 substituents independently selected from halogen, C₁₋₄ alkyl, C₃₋₈ cycloalkyl, C₁₋₆ alkyloxy, C₁₋₄ haloalkyl.

[2] The present invention also relates to compounds of formula (Ia) and (Ib) below with the variables as recited above in group [1] or [1']:



5

[3] The present invention relates to a compound as described directly above in [1], [1'] or [2] wherein

R^1 is selected from C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkylthio, $-XR^{1c}$ wherein R^1 is substituted with
 10 0-6 substituents selected from halogen, C_{1-4} alkyl or C_{1-4} haloalkyl;

R^2 is selected from C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{3-8} cycloalkyl, C_{3-6} cycloalkyl C_{1-6} alkyl, $-NR^{2c}R^{2d}$ wherein R^2 is unsubstituted or substituted with 1-3
 15 substituents independently selected from the group R^{2i} , R^{2j} , R^{2k} , C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, Br, Cl, F, I, C_{1-4} haloalkyl, $-OR^{2g}$, $-NR^{2g}R^{2h}$, $-C_{1-6}$ alkylOR^{2g}, and C_{3-8} cycloalkyl which is substituted with 0-1 R^{2i} and in which 0-1 carbons of C_{4-8} cycloalkyl is replaced by -O-.

20

[4] The present invention also relates to a compound described directly above in group [1] or [1'], [2] or [3] wherein R^3 is selected from an aryl group selected from phenyl or substituted versions thereof or a heteroaryl
 25 group selected from pyridyl or substituted versions thereof.

[5] The present invention relates to a compound described directly above in [1], [1'], [2], [3], or [4]
 30 wherein R^3 is substituted with 0-4 substituents independently selected from halogen, C_{1-4} alkyloxy, C_{1-6}

alkyl or $\text{NR}'\text{R}''$ wherein R' and R'' are independently selected from H or C_{1-6} alkyl.

[6] The present invention preferably relates to a compound as described directly above in groups [1], [1'], [2], [3], [4] or [5] wherein R^1 is selected from 2,4-dichlorophenyl, 2-chloro-4-methoxyphenyl, 2,4,6-trimethylphenyl, 2,4,6-trimethoxyphenyl, 2-dimethylamino-4-methyl-pyridin-5-yl, 2,4-dichloro-5-fluorophenyl, 2-chloro-4-methoxy-5-fluorophenyl, 2-methyl-4-methoxyphenyl, 2-methyl-4,6-dimethoxyphenyl, 2-chloro-4,5-dimethoxyphenyl or 2-chloro-4,6-dimethoxyphenyl.

[7] The present invention also preferably relates to a compound as described in the group [1], [1'], [2], [3], [4], [5], or [6] wherein R^2 is selected from C_1 alkyl of the formula $-\text{CR}'\text{R}''\text{R}'''$ wherein R' , R'' and R''' are independently selected from H, C_{1-6} alkyl, C_3 , cycloalkyl, hydroxy C_{1-6} alkyl, C_{1-6} alkyloxy C_{1-6} alkyl, C_{2-6} alkenyl, C_{1-6} alkyloxy, hydroxy, with the proviso that each of R' , R'' and R''' cannot be H;

or R^2 is selected from $\text{NR}^{2c}\text{R}^{2d}$ wherein R^{2c} and R^{2d} are independently selected from H or C_{1-6} alkyl.

[8] The present invention preferably relates to a compound according to groups [1]-[7] and [1'] wherein R^1 is selected from an aryl or heteroaryl group attached through an unsaturated carbon atom wherein, aryl is phenyl, each phenyl being substituted with 0-5 substituents independently selected at each occurrence from C_{1-6} alkyl, C_{3-6} cycloalkyl, methylenedioxy, C_{1-4} alkyloxy- C_{1-4} alkyloxy, $-\text{OR}^{2m}$, Br, Cl, F, I, C_{1-4} haloalkyl, $-\text{CN}$, $-\text{NO}_2$, $-\text{SH}$, $-\text{S}(\text{O})_n\text{R}^{2n}$, $-\text{COR}^{2m}$, $-\text{CO}_2\text{R}^{2m}$, $-\text{OC}(\text{O})\text{R}^{2n}$, $-\text{NR}^{2g}\text{COR}^{2m}$, $-\text{N}(\text{COR}^{2m})_2$, $-\text{NR}^{2g}\text{CONR}^{2c}\text{R}^{2p}$, $-\text{NR}^{2g}\text{CO}_2\text{R}^{2h}$, $-\text{NR}^{2o}\text{R}^{2p}$ and $\text{CONR}^{2o}\text{R}^{2p}$ and up to 1 phenyl, each phenyl substituent being substituted with 0-4 substituents selected from the group C_{1-3} alkyl, C_{1-3} alkoxy, Br, Cl, F, I, $-\text{CN}$, dimethylamino,


CF₃, C₂F₅, OCF₃, SO₂Me and acetyl and wherein, heteroaryl is selected at each occurrence from pyridyl, each pyridyl being substituted at 0-4 carbon atoms with a substituent independently selected at each occurrence from the group

5 C₁₋₆ alkyl, C₃₋₆ cycloalkyl, Br, F, I, C₁₋₄ haloalkyl, -CN, nitro, -OR^{2m}, -SH, -S(O)_nR²ⁿ, COR^{2m}, -CO₂R^{2m}, -OC(O)R²ⁿ, -NR^{2g}COR^{2m}, -N(COR^{2m})₂, -NR^{2g}CONR^{2o}R^{2p} and each pyridyl being substituted at any carbon atom with 0-1 substituents selected from the group R^{2g}, CO₂R^{3a}, COR^{3a} and SO₂R^{3a}.

10

[9] The present invention preferably relates to a compound of formula (Ia)

15



(Ia)

or a pharmaceutically acceptable salt thereof, wherein

20

R¹ is independently selected at each occurrence from H, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, halo, CN, C₁-C₄ haloalkyl, C₁-C₁₂ hydroxyalkyl, C₂-C₁₂ alkoxyalkyl, C₂-C₁₀ cyanoalkyl, C₃-C₆ cycloalkyl, C₄-C₁₀ cycloalkylalkyl, NR⁹R¹⁰, C₁-C₄ alkyl-NR⁹R¹⁰, NR⁹COR¹⁰, OR¹¹, SH or S(O)_nR¹²;

25

R² is selected from:

-H, OR⁷, SH, S(O)_nR¹³, COR⁷, CO₂R⁷, CHR⁶(OR⁷)R^{6a}, OC(O)R¹³, CH(OH)R⁶, C(OH)R⁶R^{6a}, C(OR⁷)R⁶R^{6a}, NO, NO₂, NR⁶COR⁷, N(COR⁷)₂, NR⁸CONR⁶R⁷, NR⁶CO₂R⁷,

30

NR^6R^7 , $\text{NR}^6\text{S}(\text{O})_2\text{R}^7$, $\text{N}(\text{S}(\text{O})_2\text{R}^7)_2$, $\text{N}(\text{OR}^7)\text{R}^6$ or CONR^6R^7 ;

or

5 -C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₃-C₈ cycloalkyl, C₅-C₈ cycloalkenyl, C₄-C₁₂ cycloalkylalkyl or C₆-C₁₀ cycloalkenylalkyl, each optionally substituted with 1 to 3 substituents independently selected at each occurrence
10 from C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄ haloalkyl, cyano, OR^{15} , SH, $\text{S}(\text{O})_n\text{R}^{13}$, COR^{15} , CO_2R^{15} , $\text{OC}(\text{O})\text{R}^{13}$, $\text{NR}^8\text{COR}^{15}$, $\text{N}(\text{COR}^{15})_2$, $\text{NR}^8\text{CONR}^{16}\text{R}^{15}$, $\text{NR}^8\text{CO}_2\text{R}^{13}$, $\text{NR}^{16}\text{R}^{15}$, $\text{CONR}^{16}\text{R}^{15}$, aryl, heteroaryl and
15 heterocyclyl;

R^3 is selected from phenyl, naphthyl, pyridyl, pyrimidinyl, triazinyl, furanyl, thienyl, benzothienyl, benzofuranyl, 2,3-dihydrobenzofuranyl, 2,3-dihydrobenzothienyl, indanyl, 1,2-benzopyranyl, 3,4-dihydro-1,2-benzopyranyl, tetralinyl, each R^3 optionally substituted with 1 to 5 substituents, each Ar is attached via an unsaturated carbon atom, wherein
20 the substituents are independently selected at each occurrence from: C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₃-C₆ cycloalkyl, C₄-C₁₂ cycloalkylalkyl, NO_2 , halo, CN, C₁-C₄ haloalkyl, NR^6R^7 , NR^8COR^7 , $\text{NR}^8\text{CO}_2\text{R}^7$, COR^7 , OR^7 , CONR^6R^7 , $\text{CO}(\text{NOR}^9)\text{R}^7$, CO_2R^7 , or $\text{S}(\text{O})_n\text{R}^7$, where each
30 such C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₃-C₆ cycloalkyl and C₄-C₁₂ cycloalkylalkyl are optionally substituted with 1 to 3 substituents independently selected at each occurrence from C₁-C₄ alkyl, NO_2 , halo, CN, NR^6R^7 , NR^6COR^7 , $\text{NR}^7\text{CO}_2\text{R}^7$, COR^7 or OR^7 , CONR^6R^7 , CO_2R^7 , $\text{CO}(\text{NOR}^9)\text{R}^7$, or $\text{S}(\text{O})_n\text{R}^7$;
35

R⁵ is selected from H, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₃-C₆ cycloalkyl, C₄-C₁₀ cycloalkylalkyl, each optionally substituted with 1 to 3 substituents independently selected at each occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl; halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹³, COR¹⁵, CO₂R¹⁵, OC(O)R¹³, NR⁸COR¹⁵, N(COR¹⁵)₂, NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹³, NR¹⁶R¹⁵, CONR¹⁶R¹⁵, aryl, heteroaryl and heterocyclyl; or halo, CN, -NR⁶R⁷, NR⁹COR¹⁰, -NR⁶S(O)_nR⁷, S(O)_nNR⁶R⁷, C₁-C₄ haloalkyl, -OR⁷, SH or -S(O)_nR¹²;

15

R⁶, R^{6a} and R⁷ are independently selected at each occurrence from:

-H,
-C₁-C₁₀ alkyl, C₃-C₁₀ alkenyl, C₃-C₁₀ alkynyl, C₁-C₁₀ haloalkyl with 1-10 halogens, C₂-C₈ alkoxyalkyl, C₃-C₆ cycloalkyl, C₄-C₁₂ cycloalkylalkyl, C₅-C₁₀ cycloalkenyl, or C₆-C₁₄ cycloalkenylalkyl, each optionally substituted with 1 to 3 substituents independently selected at each occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹³, COR¹⁵, CO₂R¹⁵, OC(O)R¹³, NR⁸COR¹⁵, N(COR¹⁵)₂, NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹³, NR¹⁶R¹⁵, CONR¹⁶R¹⁵, aryl, heteroaryl or heterocyclyl, -aryl, aryl(C₁-C₄ alkyl), heteroaryl, heteroaryl(C₁-C₄ alkyl), heterocyclyl or heterocyclyl(C₁-C₄ alkyl);

35

alternatively, NR⁶R⁷ and NR^{6a}R^{7a} are independently piperidine, pyrrolidine, piperazine, N-

methylnpiperazine, morpholine or thiomorpholine, each optionally substituted with 1-3 C₁-C₄ alkyl groups;

5 R⁸ is independently selected at each occurrence from H or C₁-C₄ alkyl;

R⁹ and R¹⁰ are independently selected at each occurrence from H, C₁-C₄ alkyl, or C₃-C₆ cycloalkyl;

10

R¹¹ is selected from H, C₁-C₄ alkyl, C₁-C₄ haloalkyl, or C₃-C₆ cycloalkyl;

R¹² is C₁-C₄ alkyl or C₁-C₄ haloalkyl;

15

R¹³ is selected from C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₂-C₈ alkoxyalkyl, C₃-C₆ cycloalkyl, C₄-C₁₂ cycloalkylalkyl, aryl, aryl(C₁-C₄ alkyl)-, heteroaryl or heteroaryl(C₁-C₄ alkyl)-;

20

R¹⁵ and R¹⁶ are independently selected at each occurrence from H, C₁-C₆ alkyl, C₃-C₁₀ cycloalkyl, C₄-C₁₆ cycloalkylalkyl, except that for S(O)_nR¹⁵, R¹⁵ cannot be H;

25

aryl is phenyl or naphthyl, each optionally substituted with 1 to 5 substituents independently selected at each occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹⁵, COR¹⁵, CO₂R¹⁵, OC(O)R¹⁵, NR⁸COR¹⁵, N(COR¹⁵)₂, NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹⁵, NR¹⁶R¹⁵, and CONR¹⁶R¹⁵;

30

heteroaryl is pyridyl, pyrimidinyl, triazinyl, furanyl, pyranyl, quinolinyl, isoquinolinyl, thienyl, imidazolyl, thiazolyl, indolyl, pyrrolyl, oxazolyl, benzofuranyl, benzothienyl, benzothiazolyl, isoxazolyl, pyrazolyl, 2,3-

35

dihydrobenzothienyl or 2,3-dihydrobenzofuranyl,
 each being optionally substituted with 1 to 5
 substituents independently selected at each
 occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl,
 5 halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹⁵,
 -COR¹⁵, CO₂R¹⁵, OC(O)R¹⁵, NR⁸COR¹⁵, N(COR¹⁵)₂,
 NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹⁵, NR¹⁶R¹⁵, and CONR¹⁶R¹⁵;

heterocyclyl is saturated or partially saturated
 10 heteroaryl, optionally substituted with 1 to 5
 substituents independently selected at each
 occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl,
 halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹⁵,
 COR¹⁵, CO₂R¹⁵, OC(O)R¹⁵, NR⁸COR¹⁵, N(COR¹⁵)₂,
 15 NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹⁵, NR¹⁵R¹⁶, and CONR¹⁶R¹⁵;

n is independently at each occurrence 0, 1 or 2.


20 [10] The present invention also relates to a compound
 according to groups [1'] and [1]-[9] wherein R³ is
 substituted with 2-4 substituents.

25 [11] The present invention also relates to a compound
 according to groups [1'] and [1]-[10] wherein R² is
 substituted with 1-4 substituents.

[12] The present invention also relates to a compound
 according to groups [1'] and [1]-[9] wherein R² is
 30 selected from 3-pentyl, NEt₂, butyl, NHCH(CH₂OMe)₂,
 NHCH(CH₂OEt)₂, NHCH(Et)CH₂OMe, NH-3-heptyl, NH-3-pentyl, NH-
 2-butyl, NH-3-hexyl, NHCH(CH₂Ph)CH₂OMe,
 NHCH(Et)CH₂CH₂OMe, NH-cyclobutyl, NH-cyclopentyl, NEtPr,
 NEtBu, NMePr, NMePh, Npr₂, NPr(CH₂-c-C₃H₅),
 35 N(CH₂CH₂OMe)₂, morpholino, N(CH₂Ph)CH₂CH₂OMe,
 N(Me)CH₂CH₂OMe, N(Et)CH₂CH₂OMe, N(CH₂-c-C₃H₅)CH₂CH₂OMe,
 N(CH₂-c-C₃H₅)Pr, N(CH₂-c-C₃H₅)Et, OEt, OCH(Et)CH₂OMe,
 OCH(Et)CH₂CH₂OMe, OCH(Me)CH₂CH₂OMe, O-3-pentyl, O-2-

- pentyl, S-3-pentyl, S-2-pentyl, SEt, S(O)Et, SO₂Et, S-3-pentyl, S(O)-3-pentyl, SO₂-3-pentyl, S-2-pentyl, S(O)-2-pentyl, SO₂-2-pentyl, CH(CO₂Et)₂, C(Et)(CO₂Et)₂, CH(Et)CH₂OH, CH(Et)CH₂OMe, CH(Et)CH₂CH₂OMe, CONMe₂,
- 5 COCH₃, COEt, COPr, CO-2-pentyl, CO-3-pentyl, CH(OH)CH₃, C(OH)Me₂, C(OH)Ph-3-pyridyl, CH(OMe)CH₃, CH(OMe)Et, CH(OMe)Pr, CH(OEt)CH₃, CH(OPr)CH₃, 2-pentyl, 2-butyl, cyclobutyl, cyclopentyl, CH(Me)cyclobutyl, CH(OMe)cyclobutyl, CH(OH)cyclobutyl, CH(Me)cyclopropyl,
- 10 CH(OMe)cyclopropyl, CH(OH)cyclopropyl, CH(Et)cyclobutyl, CH(Et)cyclopropyl, CH(OMe)cyclobutyl, CH(OMe)cyclopropyl, CH(OEt)cyclobutyl, CH(OEt)cyclopropyl, CH(Me)CH₂-cyclobutyl, CH(OMe)CH₂-cyclobutyl, CH(OH)CH₂-cyclobutyl, CH(Me)CH₂-cyclopropyl, CH(OMe)CH₂-cyclopropyl, CH(OH)CH₂-cyclopropyl,
- 15 cyclopropyl, CH(Et)CH₂-cyclobutyl, CH(Et)CH₂-cyclopropyl, CH(OMe)CH₂-cyclobutyl, CH(OMe)CH₂-cyclopropyl, CH(OEt)CH₂-cyclobutyl, CH(OEt)CH₂-cyclopropyl, CH(CH₂OMe)cyclobutyl, CH(CH₂OMe)cyclopropyl, CH(CH₂OEt)cyclobutyl, CH(CH₂OEt)cyclopropyl,
- 20 CH(cyclobutyl)₂, CH(cyclopropyl)₂, CH(Et)CH₂CONMe₂, CH(Et)CH₂CH₂NMe₂, CH(CH₂OMe)Me, CH(CH₂OMe)Et, CH(CH₂OMe)Pr, CH(CH₂OEt)Me, CH(CH₂OEt)Et, CH(CH₂OEt)Pr, CH(CH₂C≡CMe)Et, CH(CH₂C≡CMe)Et.
- 25 [13] The present invention further relates to a compound of formula I or Ia according to groups [1'] and [1]-[12] above wherein R³ is selected from 2,4-Cl₂-Ph, 2,4,6-Me₃-Ph, 2,4-Me₂-Ph, 2-Me-4-MeO-Ph, 2-Cl-4-MeO-Ph, 2-Cl-4,5-(MeO)₂-Ph, 2-Cl-4-MeO-5-F-Ph, 2-Me-4-MeO-5-F-Ph, 2,5-
- 30 (Me)₂-4-MeO-Ph, 2-Me-4-NMe₂-Ph, 2-CF₃-4-MeO-Ph, 2-Me-4-(COMe)-Ph, 2-Me-6-Me₂N-pyrid-3-yl, 4-Me-2-Me₂N-pyrid-5-yl, 2-Me-6-MeO-pyrid-3-yl, 4-Me-2-MeO-pyrid-5-yl.


- [14] The present invention also relates to a compound of
- 35 formula Ib



having R^1 - R^4 as defined in groups [1]-[13] above.

- [15] In a preferred embodiment, the present invention
 5 relates to a compound according to group [14] wherein R^1
 is selected from H, CH_3 , C_2H_5 , OCH_3 ; R^4 is selected from
 H, OCH_3 , CH_3 and C_2H_5 ; R^2 is selected from $CH(C_2H_5)_2$, $CH(c-$
 $C_3H_5)_2$, $CHC_2H_5(c-C_3H_5)$, $CH(C_2H_5)_2$, $CH(c-C_3H_5)_2$; and R^3 is
 10 2,4- Cl_2 -Ph, 2-Cl-4- CH_3O -Ph, 2,4,6- $(CH_3)_3$ -Ph,
 2-Cl-4- CF_3 -Ph and 2- $(CH_3)_2N$ -4- CH_3 -pyridin-5-yl.

[16] The present invention also relates to compounds of
 formula Ic, Id, Ie and If



15

and the pharmaceutically acceptable salts thereof wherein
 R^2 , R^3 , R^4 and R^5 are as defined in groups [1]-[15] above.

- [17] The present invention also relates to a method of
 20 antagonizing a CRF-1 receptor in mammals including humans
 wherein binding to the receptor causes and ultimately

results in the treatment of affective disorder, anxiety, depression, headache, irritable bowel syndrome, post-traumatic stress disorder, supranuclear palsy, immune suppression, Alzheimer's disease, gastrointestinal diseases, anorexia nervosa or other feeding disorder, drug addiction, drug or alcohol withdrawal symptoms, inflammatory diseases, cardiovascular or heart-related diseases, fertility problems, human immunodeficiency virus infections, hemorrhagic stress, obesity, infertility, head and spinal cord traumas, epilepsy, stroke, ulcers, amyotrophic lateral sclerosis, hypoglycemia or a disorder the treatment of which can be effected or facilitated by antagonizing CRF, including but not limited to disorders induced or facilitated by CRF, in mammals comprising administering to the mammal a therapeutically effective amount of a compound of Formula I, Ia, Ib, Ic, Id, Ie or If according to groups [1]-[16] and [1'] above with the proviso that, in the case of compounds of group [1], provisos (a) and (b) are not present.

[18] The present invention also relates to use of a compound according to groups [1]-[16] and [1'] in therapy.


[19] The present invention also provides pharmaceutical compositions comprising compounds of Formula I, Ia, Ib, Ic, Id, Ie, or If with the variables as recited above in groups [1]-[16] and [1'] and a pharmaceutically acceptable carrier.

[20] The present invention also relates to compounds according to group [9] wherein R² is NR⁶R⁷, OR⁷ or

-C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₃-C₈ cycloalkyl, C₅-C₈ cycloalkenyl, C₄-C₁₂ cycloalkylalkyl or C₆-C₁₀ cycloalkenylalkyl, each optionally

substituted with 1 to 3 substituents independently selected at each occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹³, COR¹⁵, CO₂R¹⁵, OC(O)R¹³,
5 NR⁸COR¹⁵, N(COR¹⁵)₂, NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹³, NR¹⁶R¹⁵, CONR¹⁶R¹⁵.

[21] The present invention also relates to a compound of formula II



or a pharmaceutically acceptable salt or isomer thereof wherein R¹-R⁴ and the other variables are as defined in groups [1'] and [1]-[20].

15

[22] The invention further relates to pharmaceutical compositions comprising the compound of group [21] and a pharmaceutically acceptable carrier.

20

[23] The invention also comprises use of a compound according to group [21] in therapy and to a method of treating a patient in need of treatment thereof comprising administering to said patient a pharmaceutically effective amount of a compound or

25

composition according to group [21] or [22].

Many compounds of this invention have one or more asymmetric centers or planes. Unless otherwise indicated, all chiral (enantiomeric and diastereomeric) and racemic
30 forms are included in the present invention. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds, and all such

stable isomers are contemplated in the present invention. The compounds may be isolated in optically active or racemic forms. It is well known in the art how to prepare optically active forms, such as by resolution of racemic
5 forms or by synthesis from optically active starting materials. All chiral, (enantiomeric and diastereomeric) and racemic forms and all geometric isomeric forms of a structure are intended, unless the specific stereochemistry or isomer form is specifically indicated.

10 The term "alkyl" includes both branched and straight-chain alkyl having the specified number of carbon atoms. "Alkenyl" includes hydrocarbon chains of either a straight or branched configuration and one or more unsaturated carbon-carbon bonds which may occur in
15 any stable point along the chain, such as ethenyl, propenyl, and the like. "Alkynyl" includes hydrocarbon chains of either a straight or branched configuration and one or more triple carbon-carbon bonds which may occur in any stable point along the chain, such as
20 ethynyl, propynyl and the like. "Haloalkyl" is intended to include both branched and straight-chain alkyl having the specified number of carbon atoms, substituted with 1 or more halogen; "alkoxy" represents an alkyl group of indicated number of carbon atoms
25 attached through an oxygen bridge; "cycloalkyl" is intended to include saturated ring groups, including mono-, bi- or poly-cyclic ring systems, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and so forth. "Halo" or "halogen" includes fluoro, chloro,
30 bromo, and iodo.

The term "substituted", as used herein, means that one or more hydrogen on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valency is not
35 exceeded, and that the substitution results in a stable compound. When a substituent is keto (i.e., =O), then 2 hydrogens on the atom are replaced.

Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds. By "stable compound" or "stable structure" is meant a compound that is sufficiently robust to
5 survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

The term "pharmaceutically acceptable salts" includes acid or base salts of the compounds of
10 formulas (I). Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like.

15 Pharmaceutically acceptable salts of the compounds of the invention can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of
20 the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in Remington's Pharmaceutical Sciences, 17th ed., Mack Publishing Company, Easton, PA, 1985, p. 1418, the
25 disclosure of which is hereby incorporated by reference.

"Prodrugs" are considered to be any covalently bonded carriers which release the active parent drug of formula (I) *in vivo* when such prodrug is administered
30 to a mammalian subject. Prodrugs of the compounds of formula (I) are prepared by modifying functional groups present in the compounds in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compounds.
35 Prodrugs include compounds wherein hydroxy, amine, or sulfhydryl groups are bonded to any group that, when administered to a mammalian subject, cleaves to form a free hydroxyl, amino, or sulfhydryl group,

respectively. Examples of prodrugs include, but are not limited to, acetate, formate and benzoate derivatives of alcohol and amine functional groups in the compounds of formulas (I) and the like.

5 The term "therapeutically effective amount" of a compound of this invention means an amount effective to antagonize abnormal level of CRF or treat the symptoms of affective disorder, anxiety, depression, immunological, cardiovascular or heart-related diseases
10 and colonic hypersensitivity associated with psychopathological disturbance and stress in a host.

The variables as described in group [9] above are also
15 shown in Schemes 1-4 which describe the synthesis of compounds of the invention.

20


Syntheses

Some compounds of Formula (Ia), may be prepared from
25 intermediate compounds of Formula (3) using the procedures outlined in Scheme 1 with the variables defined as above. Compounds of Formula (3) may be treated with a base in an inert solvent, followed by reaction with compounds of the Formula R^1COY , where R^1 is
30 defined above and Y is a halogen, alkoxy, dialkylamino, alkylthio, alkanoyloxy, alkanesulfonyloxy or cyano group. Bases may include, but are not limited to, alkyl lithiums, alkali metal hydrides (preferably sodium hydride), alkali metal alkoxides (1 to 6
35 carbons) (preferably sodium methoxide or sodium ethoxide), alkaline earth metal hydrides, alkali metal dialkylamides (preferably lithium di-isopropylamide), alkali metal bis(trialkylsilyl)amides (preferably sodium bis(trimethylsilyl)amide), trialkyl amines (preferably

N,N-di-isopropyl-N-ethyl amine or triethylamine) or aromatic amines (preferably pyridine). Inert solvents may include, but are not limited to, lower alkanenitriles (1 to 6 carbons, preferably acetonitrile), dialkyl ethers (preferably diethyl ether), cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), N,N-dialkylformamides (preferably dimethylformamide), N,N-dialkylacetamides (preferably

10

Scheme 1



dimethylacetamide), cyclic amides (preferably N-methylpyrrolidin-2-one), dialkylsulfoxides (preferably dimethylsulfoxide), aromatic hydrocarbons (preferably benzene or toluene) or haloalkanes of 1 to 10 carbons and 1 to 10 halogens (preferably dichloromethane). Preferred reaction temperatures range from -80°C to 100°C. The resulting intermediates (4) are then reacted with hydrazine or its hydrate in an inert solvent in the presence or absence of an acid to provide pyrazoles (5). Inert solvents may include, but are not limited to, water, alkyl alcohols (1 to 8 carbons, preferably methanol or ethanol), lower alkanenitriles (1 to 6 carbons, preferably acetonitrile), cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), N,N-dialkylformamides (preferably dimethylformamide), N,N-dialkylacetamides (preferably dimethylacetamide), cyclic amides (preferably N-methylpyrrolidin-2-one), dialkylsulfoxides (preferably dimethylsulfoxide) or aromatic hydrocarbons (preferably benzene or toluene). Acids may include, but are not limited to alkanolic acids of 2 to 10 carbons (preferably acetic acid), haloalkanoic acids (2 - 10 carbons, 1-10 halogens, such as trifluoroacetic acid), arylsulfonic acids (preferably p-toluenesulfonic acid or benzenesulfonic acid), alkanesulfonic acids of 1 to 10 carbons (preferably methanesulfonic acid), hydrochloric acid, sulfuric acid or phosphoric acid. Stoichiometric or catalytic amounts of such acids may be used. Preferred temperatures range from ambient temperature to 150°C.

Compounds of Formula (Ia) may be prepared by reaction of pyrazoles (5) with intermediates (7), or (8) or (9) in the presence or absence of an acid or base in an inert solvent. Inert solvents may include, but are not limited to, water, alkyl alcohols (1 to 8 carbons, preferably methanol or ethanol), lower alkanenitriles (1 to 6 carbons, preferably acetonitrile), cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), N,N-

dialkylformamides (preferably dimethylformamide), N,N-dialkylacetamides (preferably dimethylacetamide), cyclic amides (preferably N-methylpyrrolidin-2-one), dialkylsulfoxides (preferably dimethylsulfoxide) or

5 aromatic hydrocarbons (preferably benzene or toluene). Acids may include, but are not limited to alkanolic acids of 2 to 10 carbons (preferably acetic acid), haloalkanoic acids (2 - 10 carbons, 1-10 halogens, such as trifluoroacetic acid), arylsulfonic acids (preferably p-

10 toluenesulfonic acid or benzenesulfonic acid), alkanesulfonic acids of 1 to 10 carbons (preferably methanesulfonic acid), hydrochloric acid, sulfuric acid or phosphoric acid. Stoichiometric or catalytic amounts of such acids may be used. Bases may include, but are not

15 limited to, alkali metal carbonates, alkali metal bicarbonates, trialkyl amines (preferably N,N-diisopropyl-N-ethyl amine) or aromatic amines (preferably pyridine). Preferred temperatures range from ambient temperature to 150°C. Intermediates (7), (8) and (9) are

20 derived from amides (6). Amides (6) may be reacted in an inert solvent in the presence or absence of an acid with compounds of the Formula $R^a_2NCR^5(OR^b)_2$, where R^a and R^b independently are lower alkyl, to generate compounds of Formula (7). Inert solvents may include, but are not

25 limited to, water, alkyl alcohols (1 to 8 carbons, preferably methanol or ethanol), lower alkanenitriles (1 to 6 carbons, preferably acetonitrile), cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), N,N-dialkylformamides (preferably dimethylformamide), N,N-

30 dialkylacetamides (preferably dimethylacetamide), cyclic amides (preferably N-methylpyrrolidin-2-one), dialkylsulfoxides (preferably dimethylsulfoxide) or aromatic hydrocarbons (preferably benzene or toluene). Acids may include, but are not limited to alkanolic acids

35 of 2 to 10 carbons (preferably acetic acid), haloalkanoic acids (2 - 10 carbons, 1-10 halogens, such as trifluoroacetic acid), arylsulfonic acids (preferably p-toluenesulfonic acid or benzenesulfonic acid),


alkanesulfonic acids of 1 to 10 carbons (preferably methanesulfonic acid), hydrochloric acid, sulfuric acid or phosphoric acid. Stoichiometric or catalytic amounts of such acids may be used. Preferred temperatures range from ambient temperature to 150°C. Intermediates of Formula (7) may be converted to compounds of Formula (8) by reaction with H₂S in an inert solvent. Inert solvents may include, but are not limited to, water, alkyl alcohols (1 to 8 carbons, preferably methanol or ethanol), alkanoic acids of 2 to 10 carbons (preferably acetic acid), haloalkanoic acids (2 - 10 carbons, 1-10 halogens, such as trifluoroacetic acid), alkanesulfonic acids of 1 to 10 carbons (preferably methanesulfonic acid), hydrochloric acid, sulfuric acid, phosphoric acid or cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane). Compounds of Formula (8) may be converted to compounds of Formula (9) by treatment with a base and an alkylating agent in an inert solvent at reaction temperatures ranging from -80°C to 250°C. Bases may include, but are not limited to, alkali metal hydrides (preferably sodium hydride), alkali metal alkoxides (1 to 6 carbons) (preferably sodium methoxide or sodium ethoxide), alkaline earth metal hydrides, alkali metal dialkylamides (preferably lithium di-isopropylamide), alkali metal carbonates, alkali metal hydroxides, alkali metal bis(trialkylsilyl)amides (preferably sodium bis(trimethylsilyl)amide), trialkyl amines (preferably N,N-di-isopropyl-N-ethyl amine or triethyl amine) or aromatic amines (preferably pyridine). Alkylating agents may include, but are not limited to, C₁-C₁₀ alkyl - halides, -tosylates, -mesylates or -triflates or C₁-C₁₀ haloalkyl(1 - 10 halogens)-halides, -tosylates, -mesylates or -triflates. Inert solvents may include, but are not limited to, alkyl alcohols (1 to 8 carbons, preferably methanol or ethanol), lower alkanenitriles (1 to 6 carbons, preferably acetonitrile), dialkyl ethers (preferably diethyl ether), cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), N,N-dialkylformamides

(preferably dimethylformamide), N,N-dialkylacetamides (preferably dimethylacetamide), cyclic amides (preferably N-methylpyrrolidin-2-one), dialkylsulfoxides (preferably dimethylsulfoxide), aromatic hydrocarbons (preferably benzene or toluene) or haloalkanes of 1 to 10 carbons and 1 to 10 halogens (preferably dichloromethane). Preferred reaction temperatures range from -80°C to 100°C.

Compounds of Formula (Ia) may also be prepared by the methods shown in Scheme 2. Compounds of Formula (11) may be treated with a halogenating agent in the presence or absence of a base in the presence or absence of an inert solvent at reaction temperatures ranging from -80°C to 250°C to give products of Formula (12) (where X is halogen). Halogenating agents include, but are not limited to, Br₂, Cl₂, I₂, N-bromosuccinimide, N-iodosuccinimide or N-chlorosuccinimide. Bases may include, but are not limited to, alkali metal carbonates, alkali metal bicarbonates, trialkyl amines (preferably N,N-di-isopropyl-N-ethyl amine) or aromatic amines (preferably pyridine). Inert solvents may include, but are not limited to, lower alkanenitriles (1 to 6 carbons, preferably acetonitrile), dialkyl ethers (preferably diethyl ether), cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), N,N-dialkylformamides (preferably dimethylformamide), N,N-dialkylacetamides (preferably dimethylacetamide), cyclic amides (preferably N-methylpyrrolidin-2-one), dialkylsulfoxides (preferably dimethylsulfoxide), aromatic hydrocarbons (preferably benzene or toluene) or haloalkanes of 1 to 10 carbons and 1 to 10 halogens (preferably dichloromethane). Preferred reaction temperatures range from -20°C to 150°C. Compounds of Formula (12) may then

35

Scheme 2



- 5 be reacted with a base or a metal in the presence or
 absence of a metal salt in an inert solvent and then
 treated with a compound of the Formula R^2Y where Y is
 halogen, alkoxy, dialkylamino, alkylthio, alkanoyloxy,
 alkanesulfonyloxy or cyano groups. Examples of bases
 10 include, but are not limited to, alkyl or aryl lithiums

(e.g. n-butyl lithium or t-butyl lithium) or alkyl alkaline earth metal halides (e.g. MeMgBr). Examples of metals, include but are not limited to, alkali metals (e.g. Li) or alkali earth metals (e.g. Mg). Examples of metal salts include, but are not limited to, alkali metal halides, alkaline earth halides or transition metal halides such as ZnCl_2 , CeCl_3 or CuI . Inert solvents may include, but are not limited to dialkyl ethers (preferably diethyl ether), cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), alkanes or aromatic hydrocarbons (preferably benzene or toluene. Preferred reaction temperatures range from -100°C to 100°C .

Some compounds of Formula (Ia), where $\text{R}^2 = \text{NR}^8\text{COR}^7$, $\text{N}(\text{COR}^7)_2$, $\text{NR}^8\text{CONR}^6\text{R}^7$, $\text{NR}^8\text{CO}_2\text{R}^{13}$ or NR^6R^7 , may be prepared from intermediate compounds of Formula (11) (Formula (Ia) where $\text{R}^2 = \text{H}$), using the procedures also outlined in Scheme 2. Compounds of Formula (11) may be treated with a nitrating or nitrosating agent in the presence or absence of an acid in an inert solvent at reaction temperatures ranging from -80°C to 250°C to give products of Formula (12) (where $\text{Z} = \text{NO}$ or NO_2). Examples of nitrating agents include, but are not limited to, nitric acid, nitrous acid, alkali metal nitrates or nitrites (e.g. KNO_3 or KNO_2) or alkyl nitrites (e.g. isoamyl nitrite). Acids include, but are not limited to, alkanolic acids of 2 to 10 carbons (preferably acetic acid), haloacetic acids (e.g. trifluoroacetic acid), alkyl-, haloalkyl- or aryl-sulfonic acids (e.g. trifluoromethanesulfonic acid, p-toluenesulfonic acid or benzenesulfonic acid), alkanesulfonic acids of 1 to 10 carbons (preferably methanesulfonic acid), hydrochloric acid, sulfuric acid or phosphoric acid. Inert solvents may include, but are not limited to, water, alkyl alcohols (1 to 8 carbons, preferably methanol or ethanol), alkanes, dialkyl ethers (preferably glyme or diglyme), cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), aromatic hydrocarbons (preferably benzene or toluene). Compounds of Formula (12) may then be treated with a reducing agent in an

inert solvent to provide compounds of Formula (1), where $R^2 = NH_2$. Reducing agents include, but are not limited to, (a) hydrogen gas in combination with noble metal catalysts such as Pd-on-carbon, PtO_2 , Pt-on-carbon, Rh-on-alumina or Raney nickel or (b) alkali metal or alkaline earth metal borohydrides (preferably lithium or sodium borohydride), borane, dialkylboranes (such as di-isoamylborane), alkali metal aluminum hydrides (preferably lithium aluminum hydride), alkali metal (trialkoxo)aluminum hydrides, or dialkyl aluminum hydrides (such as di-isobutylaluminum hydride). Inert solvents may include, but are not limited to, alkyl alcohols (1 to 6 carbons), dialkyl ethers (preferably diethyl ether), cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), aromatic hydrocarbons (preferably benzene or toluene). Preferred reaction temperatures range from $-80^\circ C$ to $100^\circ C$. Compounds of Formula (Ia), where $R^2 = NH_2$, may be converted by treatment with alkylating agents or acylating agents in the presence or absence of a base in an inert solvent to compounds of Formula (Ia), where $R^2 = NR^6COR^7$, $N(COR^7)_2$, $NR^8CONR^6R^7$, $NR^6CO_2R^7$, $N(S(O)_2R^7)_2$, $NR^6S(O)_2R^7$ or NR^6R^7 . Alkylating agents may include, but are not limited to, alkyl-halides, -tosylates, -mesylates or -triflates; C_3 - C_{10} alkenyl-halides, -tosylates, -mesylates or -triflates; C_3 - C_{10} alkynyl-halides, -tosylates, -mesylates or -triflates; C_3 - C_6 cycloalkyl-halides, -tosylates, -mesylates or -triflates; C_4 - C_{12} cycloalkylalkyl-halides, -tosylates, -mesylates or -triflates; C_5 - C_{10} cycloalkenyl-halides, -tosylates, -mesylates or -triflates; or C_6 - C_{14} cycloalkenyl-halides, -tosylates, -mesylates or -triflates. Each of the above alkylating agents may be optionally substituted in a way consistent with the definition of R^2 . Acylating agents may include, but are not limited to, acyl halides or anhydrides. Sulfonylating agents include, but are not limited to, sulfonyl halides or anhydrides. Each of the above acylating or sulfonylating agents may be optionally

substituted in a way consistent with the definition of R^2 . Bases may include, but are not limited to, alkali metal hydrides (preferably sodium hydride), alkali metal alkoxides (1 to 6 carbons) (preferably sodium methoxide or sodium ethoxide), alkaline earth metal hydrides, alkali metal dialkylamides (preferably lithium diisopropylamide), alkali metal carbonates, alkali metal bis(trialkylsilyl)amides (preferably sodium bis(trimethylsilyl)amide), trialkyl amines (preferably diisopropylethyl amine) or aromatic amines (preferably pyridine). Inert solvents may include, but are not limited to, alkyl alcohols (1 to 8 carbons, preferably methanol or ethanol), lower alkanenitriles (1 to 6 carbons, preferably acetonitrile), dialkyl ethers (preferably diethyl ether), cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), N,N-dialkylformamides (preferably dimethylformamide), N,N-dialkylacetamides (preferably dimethylacetamide), cyclic amides (preferably N-methylpyrrolidin-2-one), dialkylsulfoxides (preferably dimethylsulfoxide) or aromatic hydrocarbons (preferably benzene or toluene). Preferred reaction temperatures range from $-70\text{ }^{\circ}\text{C}$ to $100\text{ }^{\circ}\text{C}$.


Compounds of Formula (Ia), where $R^2 = \text{CR}^6(\text{OR}^7)\text{R}^{6a}$, may be prepared by the procedures shown in Scheme 3.

Compounds of Formula (13) (Formula 1 where $R^3 = \text{COR}^6$) may be reacted with reagents of the Formula R^{6a}M in an inert solvent, where R^{6a} is defined above and M is alkali metal, ZnCl, ZnBr, ZnI, MgBr, MgCl, MgI, CeCl_2 , or CeBr_2 . Inert solvents may include, but are not limited to, dialkyl ethers (preferably diethyl ether), cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), or aromatic hydrocarbons (preferably benzene or toluene). Preferred reaction temperatures range from $-70\text{ }^{\circ}\text{C}$ to $100\text{ }^{\circ}\text{C}$. The resulting intermediates (15) (Formula (Ia) where $R^2 = \text{CR}^6(\text{OH})\text{R}^{6a}$) may then be reacted with an alkylating agent in the absence or presence of a base in an inert solvent. Alkylating agents may include, but are not limited to, alkyl-halides, -tosylates, -mesylates or -triflates; C_3 -

- C₁₀ alkenyl-halides, -tosylates, -mesylates or -triflates; ; C₃-C₁₀ alkynyl-halides, -tosylates, -mesylates or -triflates; C₃-C₆ cycloalkyl-halides, -tosylates, -mesylates or -triflates; C₄-
- 5 C₁₂ cycloalkylalkyl-halides, -tosylates, -mesylates or -triflates; C₅-C₁₀ cycloalkenyl-halides, -tosylates, -mesylates or -triflates; or C₆-C₁₄ cycloalkenyl-halides, -tosylates, -mesylates or -triflates. Each of the above alkylating agents may be optionally substituted in a way
- 10 consistent with the definition of R². Bases may include, but are not limited to, alkali metal hydrides (preferably sodium hydride), alkali metal alkoxides (1 to 6 carbons) (preferably sodium methoxide or sodium ethoxide), alkaline earth metal hydrides, alkali metal dialkylamides
- 15 (preferably lithium di-isopropylamide), alkali metal carbonates, alkali metal bis(trialkylsilyl)amides (preferably sodium bis(trimethylsilyl)amide), trialkyl amines (preferably di-isopropylethyl amine) or aromatic amines (preferably

20

Scheme 3




- pyridine). Inert solvents may include, but are not limited to, alkyl alcohols (1 to 8 carbons, preferably methanol or ethanol), lower alkanenitriles (1 to 6 carbons, preferably acetonitrile), dialkyl ethers (preferably diethyl ether), cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), N,N-dialkylformamides (preferably dimethylformamide), N,N-dialkylacetamides (preferably dimethylacetamide), cyclic amides (preferably N-methylpyrrolidin-2-one), dialkylsulfoxides (preferably dimethylsulfoxide) or aromatic hydrocarbons (preferably benzene or toluene). Preferred reaction temperatures range from -70 °C to 100°C.
- Alternatively, compounds of Formula (Ia), where $R^2 = CR^6(OR^7)R^{6a}$, may be prepared by other procedures shown in Scheme 3. Compounds of Formula (13) (Formula Ia where $R^2 = COR^6$) may be treated with a reducing agent in an inert.

solvent to afford intermediates of Formula (15) (Formula (Ia) where $R^2 = \text{CHR}^6\text{OH}$). Reducing agents include, but are not limited to, (a) hydrogen gas in combination with noble metal catalysts such as Pd-on-carbon, PtO_2 , Pt-on-carbon, Rh-on-alumina or Raney nickel or (b) alkali metal or alkaline earth metal borohydrides (preferably lithium or sodium borohydride), borane, dialkylboranes (such as di-isoamylborane), alkali metal aluminum hydrides (preferably lithium aluminum hydride), alkali metal (trialkoxyl)aluminum hydrides, or dialkyl aluminum hydrides (such as di-isobutylaluminum hydride). Inert solvents may include, but are not limited to, alkyl alcohols (1 to 6 carbons), dialkyl ethers (preferably diethyl ether), cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), aromatic hydrocarbons (preferably benzene or toluene). Preferred reaction temperatures range from -80°C to 100°C . Intermediates of Formula (15) (Formula (Ia) where $R^2 = \text{CHR}^6\text{OH}$) may then be reacted with an alkylating agent in the presence or absence of a base in an inert solvent. Alkylating agents may include, but are not limited to, alkyl-halides, -tosylates, -mesylates or -triflates; $\text{C}_3\text{-C}_{10}$ alkenyl-halides, -tosylates, -mesylates or -triflates; $\text{C}_3\text{-C}_{10}$ alkynyl-halides, -tosylates, -mesylates or -triflates; $\text{C}_3\text{-C}_6$ cycloalkyl-halides, -tosylates, -mesylates or -triflates; $\text{C}_4\text{-C}_{12}$ cycloalkylalkyl-halides, -tosylates, -mesylates or -triflates; $\text{C}_5\text{-C}_{10}$ cycloalkenyl-halides, -tosylates, -mesylates or -triflates; or $\text{C}_6\text{-C}_{14}$ cycloalkenyl-halides, -tosylates, -mesylates or -triflates. Each of the above alkylating agents may be optionally substituted in a way consistent with the definition of R^2 . Bases may include, but are not limited to, alkali metal hydrides (preferably sodium hydride), alkali metal alkoxides (1 to 6 carbons) (preferably sodium methoxide or sodium ethoxide), alkaline earth metal hydrides, alkali metal dialkylamides (preferably lithium di-isopropylamide), alkali metal carbonates, alkali metal bis(trialkylsilyl)amides (preferably sodium

bis(trimethylsilyl)amide), trialkyl amines (preferably di-isopropylethyl amine) or aromatic amines (preferably pyridine). Inert solvents may include, but are not limited to, alkyl alcohols (1 to 8 carbons, preferably methanol or ethanol), lower alkanenitriles (1 to 6 carbons, preferably acetonitrile), dialkyl ethers (preferably diethyl ether), cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), N,N-dialkylformamides (preferably dimethylformamide), N,N-dialkylacetamides (preferably dimethylacetamide), cyclic amides (preferably N-methylpyrrolidin-2-one), dialkylsulfoxides (preferably dimethylsulfoxide) or aromatic hydrocarbons (preferably benzene or toluene). Preferred reaction temperatures range from -70 °C to 100°C.

In addition to the specific and generic groups on compounds Ia as shown above in Schemes 1-3, the additional compounds within the generic scope having a compound of formula Ia with R¹-R⁵ as described or recited in group [1] may be made according to the general procedures described in these schemes using the appropriate starting materials and as described in the specific examples as well.


The embodiment of this invention concerning compounds of Formula (Ib) with the structure



may be prepared according to the following method:

The method of Ege and Gilbert, *J. Het. Chem.* **1981**, 18, 675-677, is used to prepare the desired ring system (Scheme 4). Thus, aminopyrazole **4-A** is converted to diazonium salt **4-B**, using sodium nitrite/acid or such reagents as isoamyl nitrite. The diazonium salt is condensed with a phosphorus ylide compound **4-C** to give the pyrazolo[5,1-c][1,2,4]triazine product.

Scheme 4



5 Some compounds of Formula (1a), may be prepared from
intermediate compounds of Formula (3) using the
procedures outlined in Scheme 5 with the variables
defined as above. Compounds of Formula (5) may be
treated with compounds of the Formula $\text{R}^5(\text{N}=\text{H})(\text{OR}')$, where
10 R' is lower alkyl or their acid- addition salts, in the
presence or absence of a base in an inert solvent. Bases
may include, but are not limited to, alkyl lithiums,
alkali metal hydrides (preferably sodium hydride), alkali
metal alkoxides (1 to 6 carbons) (preferably sodium
15 methoxide or sodium ethoxide), alkaline earth metal
hydrides, alkali metal dialkylamides (preferably lithium
di-isopropylamide), alkali metal bis(trialkylsilyl)amides
(preferably sodium bis(trimethylsilyl)amide), trialkyl
amines (preferably N,N-di-isopropyl-N-ethyl amine or
20 triethylamine) or aromatic amines (preferably pyridine).
Inert solvents may include, but are not limited to, lower
alkanenitriles (1 to 6 carbons, preferably acetonitrile),
dialkyl ethers (preferably diethyl ether), cyclic ethers
(preferably tetrahydrofuran or 1,4-dioxane), N,N-
25 dialkylformamides (preferably dimethylformamide), N,N-
dialkylacetamides (preferably dimethylacetamide), cyclic
amides (preferably N-methylpyrrolidin-2-one),

dialkylsulfoxides (preferably dimethylsulfoxide), aromatic hydrocarbons (preferably benzene or toluene) or haloalkanes of 1 to 10 carbons and 1 to 10 halogens (preferably dichloromethane). Preferred reaction

5 temperatures range from -80°C to 100°C. The resulting intermediates are then treated with compounds of the Formula R₃COY, where Y is halogen or lower alkoxy, in the presence or absence of a base in an inert solvent to provide compounds of the Formula (Ia). Bases may include,

10 but are not limited to, alkyl lithiums, alkali metal hydrides (preferably sodium hydride), alkali metal alkoxides (1 to 6 carbons) (preferably sodium methoxide or sodium ethoxide), alkaline earth metal hydrides, alkali metal dialkylamides (preferably lithium di-


15 isopropylamide), alkali metal bis(trialkylsilyl)amides (preferably sodium bis(trimethylsilyl)amide), trialkyl amines (preferably N,N-di-isopropyl-N-ethyl amine or triethylamine) or aromatic amines (preferably pyridine). Inert solvents may include, but are not limited to, lower

20 alkanenitriles (1 to 6 carbons, preferably acetonitrile), dialkyl ethers (preferably diethyl ether), cyclic ethers (preferably tetrahydrofuran or 1,4-dioxane), N,N-dialkylformamides (preferably dimethylformamide), N,N-dialkylacetamides (preferably dimethylacetamide), cyclic

25 amides (preferably N-methylpyrrolidin-2-one), dialkylsulfoxides (preferably dimethylsulfoxide), aromatic hydrocarbons (preferably benzene or toluene) or haloalkanes of 1 to 10 carbons and 1 to 10 halogens (preferably dichloromethane). Preferred reaction


30 temperatures range from -80°C to 100°C.

SCHEME 5



Compounds of formula II as described above are also readily prepared according to the general procedure described in Scheme 6. This procedure is described generally in *J. Het. Chem* **1981**, 1319.

SCHEME 6



The following specific synthetic examples describe the procedures described generically above which, when applied to appropriately substituted substrates, were and may be employed in the synthesis of the compounds in Table 1.

10

EXAMPLES

Analytical data were recorded for the compounds described below using the following general procedures. Proton NMR spectra were recorded on an Varian FT-NMR (300 MHz); chemical shifts were recorded in ppm (δ) from an internal tetramethylsilane standard in deuteriochloroform or deuterodimethylsulfoxide as specified below. Mass spectra (MS) or high resolution mass spectra (HRMS) were recorded on a Finnegan MAT 8230 spectrometer (using chemi-ionization (CI) with NH₃ as the carrier gas or gas chromatography (GC) as specified below) or a Hewlett

Packard 5988A model spectrometer. Melting points were recorded on a Buchi Model 510 melting point apparatus and are uncorrected. Boiling points are uncorrected. All pH determinations during workup were made with indicator paper.

Reagents were purchased from commercial sources and, where necessary, purified prior to use according to the general procedures outlined by D. Perrin and W.L.F. Armarego, *Purification of Laboratory Chemicals*, 3rd ed., (New York: Pergamon Press, 1988). Chromatography (thin layer (TLC) or preparative) was performed on silica gel using the solvent systems indicated below. For mixed solvent systems, the volume ratios are given. Otherwise, parts and percentages are by weight.

EXAMPLE 1 4-(2,4-dichlorophenyl)-8-(3-pentyl)-7-ethyl-2-methyl-pyrazolo[1,5-a]-1,3,5-triazine (Formula Ia, R¹ is ethyl, R⁵ is methyl, R² is 3-pentyl, R³ is 2,4-dichlorophenyl).

A. N-(1-(Dimethylamino)ethylidene)-2,4-dichlorobenzamide

A mixture of 2,4-dichlorobenzamide (1.9 g, 10 mmol) and N,N-dimethylformamide dimethyl acetal (3.5 g, 3.8 mL, 26 mmol) was heated at 120 °C and stirred under a nitrogen atmosphere for 2 h. After being cooled to room temperature, the reaction mixture was concentrated in vacuo to afford an oil. Medium pressure chromatography on silica gel (EtOAc:hexanes::1:3 to 1:1) and removal of solvent *in vacuo* afforded a solid (2.03 g, 78% yield): NMR (CDCl₃, 300 MHz): δ 7.73 (d, 1H, J = 8), 7.39 (d, 1H, J = 2), 7.24 (dd, 1H, J = 8, 2), 3.16 (s, 3H), 3.13 (s, 3H), 2.39 (s, 3H).

B. N-(Thioacetyl)-2,4-dichlorobenzamide

Hydrogen sulfide was bubbled through glacial acetic acid (20 mL) for approximately 5 min. The above

intermediate was added portionwise over 5 min. Additional hydrogen sulfide was bubbled through the reaction mixture for approximately 10 min. Nitrogen was then bubbled through the reaction mixture. Dilution with
5 water (25 mL) caused a precipitate to form. The solid was collected by filtration, washed with copious amounts of water and dried in vacuo. The resulting solid (1.75 g, 90% yield) was used without further purification: NMR (CDCl₃, 300 MHz): δ 9.9 (br s, 1H), 7.68 (d, 1H, J = 8),
10 7.51 (d, 1H, J = 2), 7.40 (dd, 1H, J = 9, 2), 3.10 (s, 3H).

C. 3-Amino-5-ethyl-4-(3-pentyl)pyrazole

15 A mixture of NaCN (4.1 g, 82.7 mmol) and KI (154 mg, 0.9 mmol) in anhydrous dimethylsulfoxide (40 mL) was heated to 40 oC with stirring. 3-Ethyl-1-bromobutane (12.4 g, 75.2 mmol) was added dropwise over 10 min. The
20 reaction mix was first heated to 80 oC and stirring was continued for 1h, then to 120 oC and stirred for 5h. The reaction mix was cooled to ambient temperature and a precipitate formed. Dilution with water (150 mL), extraction with ether (3 X 100 mL), washing the combined
25 organic layers with a saturated NaCl solution, drying over MgSO₄ and filtration afforded a solution. Removal of solvent in vacuo provided 3-ethylpentanenitrile (7.4 g, 89% yield): NMR (CDCl₃, 300 MHz): δ 2.33 (d, 2H, J = 6), 1.45-1.40 (m, 5H), 0.92 (t, 6H, J = 7).

30 A solution of di-isopropylamine (14.8 g, 20.5 mL, 147 mmol) in anhydrous THF (40 mL) was cooled to -78 oC with stirring under a nitrogen atmosphere. A solution of n-butyl lithium in hexanes (1.6M, 87.5 mL, 140 mmol) was added dropwise over 15 min. The resulting solution was stirred for an additional 30 min. A solution of 3-
35 ethylpentanenitrile (7.4 g, 67 mmol) in THF (30 mL) was added dropwise over 15 min and then the reaction mixture was stirred for 30 min. A solution of ethyl propionate (6.8 g, 7.6 mL, 67 mmol) in THF (20 mL) was added

dropwise; then the reaction mixture was warmed with stirring to ambient temperature over 3 h. The mix was poured onto water (200 mL) and the pH was adjusted to ~ 4 by the slow addition of a concentrated HCl solution.

- 5 Three extractions with ether (100 mL), drying the combined organic layers over MgSO₄, filtration and removal of solvent in vacuo provided an oil (9.53 g): NMR (CDCl₃, 300 MHz): δ 3.50 (d, 1H, J = 4), 2.74 (q, 2H, J = 7), 1.85-1.75 (m, 1H), 1.45-1.35 (m, 4H), 1.12 (t, 3H, J = 7), 1.0-0.8 (m, 6H).

- 10 A mixture of the above intermediate (7.0 g), hydrazine hydrate (2.30 g, 2.23 mL, 46 mmol) and glacial acetic acid (1 mL) in toluene was heated to reflux temperature in a Dean-Stark apparatus and stirred for 16 h. The reaction mixture was cooled to ambient temperature and solvent was removed in vacuo. EtOAc (100 mL) was added to the residue and the resulting solution was washed three times with a saturated NaHCO₃ solution (25 mL). The organic solution was dried over MgSO₄,
20 filtered and concentrated in vacuo to provide an oil (6.7 g, 88% overall yield): NMR (CDCl₃, 300 MHz): δ 2.53 (q, 2H, J = 7), 1.70-1.50 (m, 5H), 1.20 (t, 3H, J = 7), 0.83 (t, 6H, J = 7).

- 25 D. A mixture of N-(thioacetyl)-2,4-dichlorobenzamide (200 mg, 0.81 mmol) and 3-amino-5-ethyl-4-(3-pentyl)pyrazole (146 mg, 0.81 mmol) in dioxan (1 mL) was stirred at reflux temperature for 16 h. After being cooled to room temperature, the reaction mixture was concentrated in
30 vacuo. Preparative TLC using EtOAc:hexanes::1:1 generated the title product (77.3 mg, 24% yield): mp = 91-93 °C; NMR (CDCl₃, 300 MHz): δ 7.64 (d, 1H, J = 8), 7.57 (d, 1H, J = 2), 7.44 (dd, 1H, J = 8, 2), 2.77 (q, 2H, J = 8), 2.69 (s, 3H), 2.80-2.60 (m, 1H), 2.00-1.80 (m, 4H),
35 1.25 (t, 3H, J = 8), 0.82 (t, 6H, J = 7); CI-HRMS m/z Calcd: 377.1294, Found: 377.1303.

EXAMPLE 1a 4-(2,4-dichlorophenyl)-8-(3-pentyl)-7-ethyl-2-methyl-pyrazolo[1,5-a]-1,3,5-triazine

Alternate Preparation of (Formula Ia, R¹ is ethyl, R⁵ is methyl, R² is 3-pentyl, R³ is 2,4-dichlorophenyl)

5

A. N-(1-(Methylthio)ethylidene)-2,4-dichlorobenzamide

A mixture of N-(thioacetyl)-2,4-dichlorobenzamide (400 mg, 1.6 mmol) and K₂CO₃ (445 mg, 3.22 mmol) in anhydrous acetonitrile (20 mL) was stirred at ambient temperature. Iodomethane (458 mg, 0.2 mL, 3.22 mmol) was added and the reaction mixture was then stirred for 2.5 h. Solvent was removed in vacuo and the residue was partitioned between ether and water. The combined organic layers were dried over MgSO₄, filtered and concentrated in vacuo to give an oil (406 mg): NMR (CDCl₃, 300 MHz): δ 7.86 (d, 1H, J = 8), 7.47 (d, 1H, J = 2), 7.32 (dd, 1H, J = 8, 2), 2.40 (s, 3H), 2.29 (s, 3H).

B. A mixture of N-(1-(methylthio)ethylidene)-2,4-dichlorobenzamide (100 mg, 0.38 mmol) and 3-amino-5-ethyl-4-(3-pentyl)pyrazole (69 mg, 0.38 mmol) in dioxan (1 mL) was stirred at reflux temperature for 2 h. After being cooled to room temperature, the reaction mixture was concentrated in vacuo. Preparative TLC using EtOAc:hexanes::1:1 generated the title product (68 mg, 47% yield), which was identical to the product obtained by the method described in Example 1.

30

EXAMPLE 2 4-(2,4-dichlorophenyl)-8-diethylamino-7-ethyl-2-methyl-pyrazolo[1,5-a]-1,3,5-triazine

(Formula 1, R¹ is ethyl, R⁵ is methyl, R² is diethylamino, R³ is 2,4-dichlorophenyl)

35

A. 4-(2,4-dichlorophenyl)-7-ethyl-2-methyl-pyrazolo[1,5-a]-1,3,5-triazine

A mixture of 3-amino-5-ethylpyrazole (10.3 g, 39.3 mmol) and N-(1-(methylthio)ethylidene)-2,4-dichlorobenzamide (4.0 g, 35.7 mmol) in anhydrous dioxan (20 mL) was stirred at reflux temperature under a nitrogen atmosphere for 16 h. After being cooled to ambient temperature, the reaction mix was concentrated in vacuo and the residue was treated with dichloromethane. The resulting suspension was filtered and the filtrate was concentrated in vacuo to afford an oil (1.5 g, 14 % yield): NMR (CDCl₃, 300 MHz): δ 7.62 (d, 1H, J = 8), 7.59 (d, 1H, J = 2), 7.45 (dd, 1H, J = 8, 2), 6.42 (s, 1H), 2.82 (q, 2H, J = 8), 2.73 (s, 3H), 1.30 (t, 3H, J = 8).

B. 4-(2,4-dichlorophenyl)-7-ethyl-2-methyl-8-nitropyrazolo[1,5-a]-1,3,5-triazine

A solution of 4-(2,4-dichlorophenyl)-7-ethyl-2-methyl-pyrazolo[1,5-a]-1,3,5-triazine (653 mg, 2.2 mmol) in acetic anhydride (4 mL) was cooled with stirring to -5 to -10 °C. A solution of fuming nitric acid (160 mg, 2.55 mmol) in acetic anhydride (2 mL) was added dropwise over 15 min. The reaction mixture was stirred at the same temperature for 3 h, then it was partitioned between water and ether three times. The combined organic layers were washed with a saturated NaHCO₃ solution, dried over MgSO₄ and filtered. Solvent was removed in vacuo to provide a solid. Column chromatography (EtOAc:hexanes::1:9 to 1:4) provided a solid (166 mg, 21% yield) after removal of solvent in vacuo: NMR (CDCl₃, 300 MHz): δ 7.64 (d, 1H, J = 8), 7.63 (d, 1H, J = 2), 7.50 (dd, 1H, J = 8, 2), 3.19 (q, 2H, J = 7), 2.94 (s, 3H), 1.32 (t, 3H, J = 7).

B. 4-(2,4-dichlorophenyl)-7-ethyl-2-methyl-8-aminopyrazolo[1,5-a]-1,3,5-triazine

A mixture of the above intermediate (160 mg, 0.45 mmol), Na₂S₂O₄ (554 mg, 3.2 mmol), concentrated ammonium

hydroxide (0.1 mL) in dioxan (8 mL) was stirred at ambient temperature for 30 min. The reaction mixture was concentrated in vacuo and the residue was partitioned between ether and water three times. The organic layers
5 were combined, dried over MgSO₄, filtered and concentrated in vacuo to give a solid (92.5 mg, 64% yield): NMR (CDCl₃, 300 MHz): δ 7.60 (d, 1H, J = 8), 7.57 (d, 1H, J = 2), 7.43 (dd, 1H, J = 8, 2), 2.79 (q, 2H, J = 8), 2.63 (s, 3H), 1.30 (t, 3H, J = 8)

10

C.

A mixture of the above intermediate (46 mg, 0.14 mmol), ethyl triflate (64 mg, 46 μ L, 0.36 mmol) and i-Pr₂NEt (46 mg, 62 μ L, 0.36 mmol) in dichloromethane (1
15 mL) was stirred at room temperature for 2h. Solvent was removed in vacuo. Column chromatography (EtOAc:hexanes::1:9) generated the title compound, a solid (30.1 mg, 57% yield): mp = 98-99 °C: NMR (CDCl₃, 300 MHz): δ 7.63 (d, 1H, J = 8), 7.57 (d, 1H, J = 2), 7.44
20 (dd, 1H, J = 8, 2), 3.22 (q, 4H, J = 7), 2.77 (q, 2H, J = 8), 2.69 (s, 3H), 1.25 (t, 3H, J = 8), 1.00 (t, 6H, J = 8); CI-HRMS m/z Calcd: 378.1252; Found: 378.1274.

Using the above procedures and modifications known
25 to one skilled in the art of organic synthesis, the following examples of Table 1 were or may be prepared. The examples delineated in Table 1 may be prepared by the methods outlined in Examples 1, 2 or 3 or combinations thereof. Commonly used abbreviations are: Ph is phenyl,
30 Pr is propyl, Me is methyl, Et is ethyl, Bu is butyl, Ex is Example, amorph. is amorphous. In Table 1, unless otherwise indicated, the examples with the physical data shown are based upon structure Ic. Example 10, described below, shows a detailed preparation of a 1,2,4 triazine.
35 Table 1 also shows the preferred examples having structures Ie and If that can readily be made according to the procedure delineated below for Example 10.

Example 10 4-(2,4-dichlorophenyl)-7-ethyl-8-(3-pentyl)pyrazolo[5,1-c][1,2,4]triazine

Part A. A mixture of 2-ethyl-1-bromobutane (10.0 mL, 71.4 mmol), potassium cyanide (14.0 g, 215 mmol) and aliquat 336 (10 drops) in 50 mL water was heated to reflux overnight with vigorous stirring. The mixture was cooled, and extracted with dichloromethane (2 x 50 mL). The extracts were combined, dried over magnesium sulfate, filtered and evaporated. The residual liquid was distilled bulb-to-bulb to afford pure product, 3-ethylpentanenitrile (5.50 g, 49.5 mmol, 69%). b.p. 40-45 °C (5 mm Hg). Spectral data: ¹H NMR (300 MHz, CDCl₃): δ 2.33 (2H, d, J = 5.8 Hz), 1.62-1.36 (5H, m), 0.92 (6H, t, J = 7.3 Hz). MS (H₂O-GC/MS): m/e 112 (100).

Part B. A solution of diisopropylamine (7.50 mL, 57.2 mmol) in THF (100 mL) was cooled to -78 °C, and treated with n-butyllithium (34.0 mL of a 1.6 M solution in hexane). The solution was warmed briefly to 0 °C, and then recooled to -78 °C. The nitrile compound from Part A was then added by syringe, and the solution was allowed to stir for 1 hour. Then, ethyl propionate (6.50 mL, 56.7 mmol) was added by syringe, and the resulting mixture was allowed to stir and warm to ambient temperature for 12 hours. It was poured into 200 mL of satd. aq. NH₄Cl solution, and this was extracted with ethyl acetate (2 x 200 mL). The extracts were combined, dried over magnesium sulfate, filtered and evaporated. The residual oil was separated by column chromatography (silica gel, 10:90 ethyl acetate-hexane) to afford the product, 4-cyano-5-ethyl-3-heptanone, as an oil 4.06 g, 24.3 mmol, 49%). TLC R_f 0.47 (20:80 ethyl acetate-hexane). Spectral data: ¹H NMR (300 MHz, CDCl₃): δ 3.49 (1H, d, J = 4.4 Hz), 2.74 (2H, q, J = 7.3 Hz), 2.08-1.98 (1H, m), 1.70-1.58 (1H, m), 1.50-1.20 (3H, m), 1.12 (3H, t, J = 7.3 Hz), 0.95 (3H, t, J = 7.3

Hz), 0.91 (3H, t, J = 7.3 Hz). MS (H₂O-GC/MS): m/e 167 (100).

Part C. A solution of the ketonitrile from Part B (4.06 g, 24.3 mmol), hydrazine hydrate (2.70 mL, 55.7 mmol) and acetic acid (5.00 mL, 83.7 mmol) in benzene (50 mL) was heated to reflux under a Dean-Stark trap with azeotropic distillation of water. After being heated for 12 hours, the mixture was cooled and poured into 100 mL 1 N aq. NaHCO₃ solution. This was extracted with ethyl acetate (2 x 100 mL), and the extracts were washed in sequence with brine, combined, dried over sodium sulfate, filtered and evaporated to afford sufficiently-pure product, 3-amino-5-ethyl-4-(3-pentyl)pyrazole, as a viscous oil (2.48 g, 13.7 mmol, 56%). Spectral data: ¹H NMR (300 MHz, CDCl₃): δ 3.48 (2H, br), 2.54 (2H, q, J = 7.3 Hz), 2.25-2.14 (1H, m), 1.71-1.49 (4H, m), 1.20 (3H, t, J = 7.3 Hz), 0.83 (6H, t, J = 7.3 Hz), 1H missing. MS (NH₃-CI): m/e 183 (12), 182 (100).

Part D. A solution of 3-amino-5-ethyl-4-(3-pentyl)pyrazole (0.750 g, 4.14 mmol) was suspended in 4 mL water and made acidic with conc. aq. HCl (2 mL). This was cooled in an ice bath, and a conc. aq. solution of sodium nitrite (0.286 g, 4.14 mmol) was added dropwise. After stirring for 30 min., the mixture was diluted with ice-cold dichloromethane (40 mL) and made alkaline with a saturated solution of sodium carbonate in water. The organic layer was separated, dried over sodium sulfate and filtered. The filtrate was then delivered dropwise to a stirring solution of 2,4-dichlorobenzoyl)methylenetriphenylphosphorane (Bauer, et al., *J. Het. Chem.* **1998**, 35, 81-87) (1.86 g, 4.14 mmol) in dichloromethane at 10 °C. After stirring for 10 hrs. with warming to ambient temperature, the reaction mixture was evaporated, and the residual material was separated by column chromatography (silica gel, 15:85 ethyl acetate-hexane) to afford the title product (0.124 g,

8.2%) as orange crystals (m.p. 116.7-117.8 °C). TLC R_f 0.51 (20:80 ethyl acetate-hexane).

This procedure can be utilized to make compounds of formula Ib having the variables as defined in group [1] above by appropriately substituting or preparing any of the different variables for R^1 , R^2 , R^3 , R^4 including substituted versions thereof as defined in group [1] or any of the more preferred groups.

Example 11

4-phenyl-8-(3-pentyl)-7-ethyl-2-methyl-pyrazolo[1,5-a]-1,3,5-triazine (Formula Ia, R^1 is ethyl, R^5 is methyl, R^2 is 3-pentyl, R^3 is phenyl).

Ethyl acetimidate hydrochloride (656 mg, 5.31 mmol) was added to a solution of K_2CO_3 (734 mg, 5.31 mmol) in H_2O (2 mL) in a separatory funnel. The aqueous layer was extracted with CH_2Cl_2 (3 x 1 mL) to form the free base. The combined organic layers were dried with Na_2SO_4 , and filtered through a plug of cotton. The CH_2Cl_2 extract was transferred directly into a 15 mL round bottom flask containing of 3-amino-5-ethyl-4-(3-pentyl)pyrazole (300 mg, 1.77 mmol). Acetonitrile (1.5 mL, anhydrous) was added followed by HOAc (0.112 mL, 1.95 mmol) and the mixture was stirred overnight at room temperature. The solid was collected by filtration to give 358 mg (96% yield) of 3-acetamidino-5-ethyl-4-(3-pentyl)pyrazole, hydrochloride salt as a white solid: mp = 168.5-171.5 °C, 1H NMR, 300 MHz (D_2O) δ 2.51 (q, J = 7.7 Hz, 2 H), 2.24 (s, 3 H), 2.21 - 2.15 (m, 1 H), 1.72 (s, 3 H), 1.52 - 1.43 (m, 2 H), 1.33 - 1.22 (m, 2 H), 1.02 (t, J = 7.7 Hz, 3 H), 0.56 (t, J = 7.3 Hz, 6 H), CI-MS (NH_3) m/e 223.2 [(M + H - HOAc) $^+$; calcd for $C_{12}H_{23}N_4$: 223.2].

To a solution of K_2CO_3 (48 mg, 0.35 mmol) in H_2O (3 mL) in a separatory funnel was added 3-acetamidino-5-ethyl-4-(3-pentyl)pyrazole, hydrochloride salt (60 mg, 0.212 mmol). The aqueous layer was extracted with CH_2Cl_2 (4 x 5 mL). The combined organic layers were washed with brine, dried over Na_2SO_4 , filtered, and concentrated to give the free base of the pyrazole (39 mg, 0.175 mmol).

This intermediate (39 mg, 0.175 mmol) was dissolved in dioxane (1 mL) and a solution of benzoyl chloride (30 mg, 0.210 mmol) in dioxane (1 mL) was added via cannula followed by the addition of a catalytic amount of 4-
5 dimethylaminopyridine. The mixture was stirred at room temperature for 15 min (turned cloudy then clear) and was subsequently heated at reflux (112 °C) for 15 h. The mixture was cooled to room temperature and concentrated. The residue was purified via preparative thin layer
10 chromatography (silica gel, 1 mm thickness) using 20% EtOAc in hexanes to give 29 mg (54% yield) of the title compound as a yellow oil: $R_f = 0.66$, $^1\text{H NMR}$, 300 MHz (CDCl₃) δ 8.75 - 8.71 (m, 2 H), 7.64 - 7.53 (m, 3 H), 2.85 (q, $J = 7.7$ Hz, 2 H), 2.72 - 2.62 (m, 1 H), 2.70 (s, 3
15 H), 1.97 - 1.76 (m, 4 H), 1.37 (t, $J = 7.6$ Hz, 3 H), 0.82 (t, $J = 7.3$ Hz, 6 H), LRMS (CI, NH₃) m/e 309.2 [(M + H)⁺; calcd for C₁₉H₂₅N₄: 309.2].

Example 12

20 4-phenyl-8-(3-pentyl)-7-ethyl-2-methyl-pyrazolo[1,5-a]-1,3,5-triazine (Formula Ia, R¹ is ethyl, R⁵ is methyl, R² is 3-pentyl, R³ is 2-methyl-4-chlorophenyl).

To a solution of K₂CO₃ (48 mg, 0.35 mmol) in H₂O (3
25 mL) in a separatory funnel was added 3-acetamido-5-ethyl-4-(3-pentyl)pyrazole, hydrochloride salt (49 mg, 0.174 mmol). The aqueous layer was extracted with CH₂Cl₂ (4 x 5 mL). The combined organic layers were washed with brine, dried over Na₂SO₄, filtered, and concentrated to
30 give the free base of **2** (31 mg, 0.139 mmol). This intermediate was dissolved in dioxane (1 mL) and a solution of 4-chloro-2-methylbenzoyl chloride (32 mg, 0.167 mmol) in dioxane (1 mL) was added via cannula followed by the addition of a catalytic amount of 4-
35 dimethylaminopyridine. The mixture was stirred at room temperature for 15 min (turned cloudy then clear) and was subsequently heated at reflux (112 °C) for 15 h. The

mixture was cooled to room temperature and concentrated. The residue was purified via preparative thin layer chromatography (silica gel, 1 mm thickness) using 10% EtOAc in hexanes to give 14 mg (28% yield) of the title compound as a yellow solid: $R_f = 0.40$, mp = 84.5 - 86.5 °C, ^1H NMR, 300 MHz (CDCl_3) δ 7.64 (d, $J = 8.1$ Hz, 1 H), 7.35 (s, 1 H), 7.34 (d, $J = 7.7$ Hz, 1 H), 2.77 (q, $J = 7.7$ Hz, 2 H), 2.68 (s, 3 H), 2.67 - 2.60 (m, 1 H), 2.28 (s, 3 H), 1.97 (m, 4 H), 1.25 (t, $J = 7.7$ Hz, 3 H), 0.82 (t, $J = 7.3$ Hz, 6 H), CI-MS (NH_3) m/e 357.1 [$(\text{M} + \text{H})^+$; calcd for $\text{C}_{26}\text{H}_{26}\text{N}_4\text{Cl}$: 357.2].


Example 13

4-(2-chloro-4-methylsulfonylphenyl)-8-(3-pentyl)-7-ethyl-2-methyl-pyrazolo[1,5-a]-1,3,5-triazine (Formula Ia, R^1 is ethyl, R^5 is methyl, R^2 is 3-pentyl, R^3 is 2-methyl-4-methylsulfonylphenyl).

To a solution of K_2CO_3 (280 mg, 2.03 mmol) in H_2O (15 mL) in a separatory funnel was added 3-acetamidino-5-ethyl-4-(3-pentyl)pyrazole, hydrochloride salt (250 mg, 1.12 mmol). The aqueous layer was extracted with CH_2Cl_2 (4 x 15 mL). The combined organic layers were washed with brine, dried over Na_2SO_4 , filtered, and concentrated to give the free base of 3-acetamidino-5-ethyl-4-(3-pentyl)pyrazole, hydrochloride salt (225 mg, 1.01 mmol). This intermediate was dissolved in dioxane (2 mL) and a solution of 2-chloro-4-methanesulfonylbenzoyl chloride (307 mg, 1.21 mmol) in dioxane (2 mL) was added via cannula followed by the addition of DMAP (cat.). The mixture was stirred at room temperature for 15 min (turned cloudy then clear) and was subsequently heated at reflux (112 °C) for 15 h. The mixture was cooled to room temperature and concentrated. The residue was purified via prep plate using 50% EtOAc in hexanes to give 95 mg (22% yield) of the title compound as a yellow solid: $R_f = 0.61$, mp = 172.3 - 173.8 °C, ^1H NMR, 300 MHz (CDCl_3) δ 8.14 (d, $J = 1.4$ Hz, 1 H), 8.03 (dd, $J = 6.6, 1.5$ Hz, 1

H), 7.91 (d, $J = 8.0$ Hz, 1 H), 3.13 (s, 3 H), 2.78 (q, $J = 7.6$ Hz, 2 H), 2.70 (s, 3 H), 2.71 - 2.59 (m, 1 H), 2.00-1.77 (m, 4 H), 1.25 (t, $J = 7.5$ Hz, 3 H), 0.83 (t, $J = 7.5$ Hz, 6 H), ESI-MS m/e 421.0 [(M + H) $^+$]; calcd for
5 C₂₀H₂₆N₄O₂SCl: 421.1.

TABLE 1



| 5 | Ex. | Formula | R^5 | R^4 | R^2 | R^1 | mp (°C) |
|----|-----|---------|-------|-------|--|---------------------------------|---------|
| | 1 | Ic | Me | - | 3-pentyl | 2,4-Cl ₂ -Ph | 91-93 |
| | 3 | Ic | Me | - | NEt ₂ | 2,4-Cl ₂ -Ph | 98-99 |
| | 4 | Ic | Me | - | 3-pentyl | 2-Me-4-MeO-Ph | 65-67 |
| 10 | 5 | Ic | Me | - | 3-pentyl | 2-Cl-4,5-(MeO) ₂ -Ph | 104-105 |
| | 6 | Ic | Me | - | 3-pentyl | 2-Cl-4-MeO-5-F-Ph | 94-97 |
| | 7 | Ic | Me | - | 3-pentyl | 2-Me-4-MeO-5-F-Ph | 105-107 |
| | 8 | Ic | Me | - | 3-pentyl | 2,6-Me ₂ -pyrid-3-yl | oil |
| | 9 | Ic | Me | - | butyl | 2,4-Cl ₂ -Ph | amorph. |
| 15 | 10 | Ie | - | H | 3-pentyl | 2,4-Cl ₂ -Ph | |
| | 11 | Ic | Me | H | NHCH(CH ₂ OMe) ₂ | 2,4-Cl ₂ -Ph | |
| | 12 | Ic | Me | H | NHCH(Et)CH ₂ OMe | 2,4-Cl ₂ -Ph | |

| | | | | | | |
|----|----|----|----|---|--|---------------------------|
| | 13 | Ic | Me | H | NH-2-butyl | 2,4-Cl ₂ -Ph |
| | 14 | Ic | Me | H | OCH(Et)CH ₂ OMe | 2,4-Cl ₂ -Ph |
| | 15 | Ic | Me | H | O-3-pentyl | 2,4-Cl ₂ -Ph |
| | 16 | Ic | Me | H | O-2-pentyl | 2,4-Cl ₂ -Ph |
| 5 | 17 | Ic | Me | H | R-2-pentyl | 2,4-Cl ₂ -Ph |
| | 18 | Ic | Me | H | S-2-pentyl | 2,4-Cl ₂ -Ph |
| | 19 | Ic | Me | H | R-2-butyl | 2,4-Cl ₂ -Ph |
| | 20 | Ic | Me | H | S-2-butyl | 2,4-Cl ₂ -Ph |
| | 21 | Ic | Me | H | CH(Et)CH ₂ OH | 2,4-Cl ₂ -Ph |
| 10 | 22 | Ic | Me | H | CH(Et)CH ₂ OMe | 2,4-Cl ₂ -Ph |
| | 23 | Ic | Me | H | COCH ₃ | 2,4-Cl ₂ -Ph |
| | 24 | Ic | Me | H | COEt | 2,4-Cl ₂ -Ph |
| | 25 | Ic | Me | H | CO ₂ Et | 2,4-Cl ₂ -Ph |
| | 26 | Ic | Me | H | CO-2-pentyl | 2,4-Cl ₂ -Ph |
| 15 | 27 | Ic | Me | H | CO-3-pentyl | 2,4-Cl ₂ -Ph |
| | 28 | Ic | Me | H | CH(OH)CH ₃ | 2,4-Cl ₂ -Ph |
| | 29 | Ic | Me | H | C(OH)Me ₂ | 2,4-Cl ₂ -Ph |
| | 30 | Ic | Me | H | C(OH)Ph-3-pyridyl | 2,4-Cl ₂ -Ph |
| | 31 | Ic | Me | H | CH(OMe)CH ₃ | 2,4-Cl ₂ -Ph |
| 20 | 32 | Ic | Me | H | CH(OMe)Et | 2,4-Cl ₂ -Ph |
| | 33 | Ic | Me | H | CH(OMe)Pr | 2,4-Cl ₂ -Ph |
| | 34 | Ic | Me | H | CH(OEt)CH ₃ | 2,4-Cl ₂ -Ph |
| | 35 | Ic | Me | H | CH(OPr)CH ₃ | 2,4-Cl ₂ -Ph |
| | 36 | Ic | Me | H | CH(OMe)Et | 2,4-Cl ₂ -Ph |
| 25 | 37 | Ic | Me | H | CH(OMe)Pr | 2,4-Cl ₂ -Ph |
| | 38 | Ic | Me | H | cyclobutyl | 2,4-Cl ₂ -Ph |
| | 39 | Ic | Me | H | cyclopentyl | 2,4-Cl ₂ -Ph |
| | 40 | Ic | Me | H | CH(Me)cyclobutyl | 2,4-Cl ₂ -Ph |
| | 41 | Ic | Me | H | CH(OMe)cyclobutyl | 2,4-Cl ₂ -Ph |
| 30 | 42 | Ic | Me | H | CH(Me)cyclopropyl | 2,4-Cl ₂ -Ph |
| | 43 | Ic | Me | H | CH(OMe)cyclopropyl | 2,4-Cl ₂ -Ph |
| | 44 | Ic | Me | H | CH(Et)cyclobutyl | 2,4-Cl ₂ -Ph |
| | 45 | Ic | Me | H | CH(OEt)cyclobutyl | 2,4-Cl ₂ -Ph |
| | 46 | Ic | Me | H | CH(Et)cyclopropyl | 2,4-Cl ₂ -Ph |
| 35 | 47 | Ic | Me | H | CH(OEt)cyclopropyl | 2,4-Cl ₂ -Ph |
| | 48 | Ic | Me | H | CH(cyclobutyl) ₂ | 2,4-Cl ₂ -Ph |
| | 49 | Ic | Me | H | CH(cyclopropyl) ₂ | 2,4-Cl ₂ -Ph |
| | 50 | Ic | Me | H | NHCH(CH ₂ OMe) ₂ | 2,4,6-Me ₃ -Ph |

| | | | | | | |
|----|----|----|----|---|-----------------------------|---------------------------|
| | 51 | Ic | Me | H | NHCH(Et)CH ₂ OMe | 2,4,6-Me ₃ -Ph |
| | 52 | Ic | Me | H | NH-2-butyl | 2,4,6-Me ₃ -Ph |
| | 53 | Ic | Me | H | OCH(Et)CH ₂ OMe | 2,4,6-Me ₃ -Ph |
| | 54 | Ic | Me | H | O-3-pentyl | 2,4,6-Me ₃ -Ph |
| 5 | 55 | Ic | Me | H | O-2-pentyl | 2,4,6-Me ₃ -Ph |
| | 56 | Ic | Me | H | R-2-pentyl | 2,4,6-Me ₃ -Ph |
| | 57 | Ic | Me | H | S-2-pentyl | 2,4,6-Me ₃ -Ph |
| | 58 | Ic | Me | H | R-2-butyl | 2,4,6-Me ₃ -Ph |
| | 59 | Ic | Me | H | S-2-butyl | 2,4,6-Me ₃ -Ph |
| 10 | 60 | Ic | Me | H | 3-pentyl | 2,4,6-Me ₃ -Ph |
| | 61 | Ic | Me | H | CH(Et)CH ₂ OH | 2,4,6-Me ₃ -Ph |
| | 62 | Ic | Me | H | CH(Et)CH ₂ OMe | 2,4,6-Me ₃ -Ph |
| | 63 | Ic | Me | H | COCH ₃ | 2,4,6-Me ₃ -Ph |
| | 64 | Ic | Me | H | COEt | 2,4,6-Me ₃ -Ph |
| 15 | 65 | Ic | Me | H | CO ₂ Et | 2,4,6-Me ₃ -Ph |
| | 66 | Ic | Me | H | CO-2-pentyl | 2,4,6-Me ₃ -Ph |
| | 67 | Ic | Me | H | CO-3-pentyl | 2,4,6-Me ₃ -Ph |
| | 68 | Ic | Me | H | CH(OH)CH ₃ | 2,4,6-Me ₃ -Ph |
| | 69 | Ic | Me | H | C(OH)Me ₂ | 2,4,6-Me ₃ -Ph |
| 20 | 70 | Ic | Me | H | C(OH)Ph-3-pyridyl | 2,4,6-Me ₃ -Ph |
| | 71 | Ic | Me | H | CH(OMe)CH ₃ | 2,4,6-Me ₃ -Ph |
| | 72 | Ic | Me | H | CH(OMe)Et | 2,4,6-Me ₃ -Ph |
| | 73 | Ic | Me | H | CH(OMe)Pr | 2,4,6-Me ₃ -Ph |
| | 74 | Ic | Me | H | CH(OEt)CH ₃ | 2,4,6-Me ₃ -Ph |
| 25 | 75 | Ic | Me | H | CH(OPr)CH ₃ | 2,4,6-Me ₃ -Ph |
| | 76 | Ic | Me | H | CH(OMe)Et | 2,4,6-Me ₃ -Ph |
| | 77 | Ic | Me | H | CH(OMe)Pr | 2,4,6-Me ₃ -Ph |
| | 78 | Ic | Me | H | cyclobutyl | 2,4,6-Me ₃ -Ph |
| | 79 | Ic | Me | H | cyclopentyl | 2,4,6-Me ₃ -Ph |
| 30 | 80 | Ic | Me | H | CH(Me)cyclobutyl | 2,4,6-Me ₃ -Ph |
| | 81 | Ic | Me | H | CH(OMe)cyclobutyl | 2,4,6-Me ₃ -Ph |
| | 82 | Ic | Me | H | CH(Me)cyclopropyl | 2,4,6-Me ₃ -Ph |
| | 83 | Ic | Me | H | CH(OMe)cyclopropyl | 2,4,6-Me ₃ -Ph |
| | 84 | Ic | Me | H | CH(Et)cyclobutyl | 2,4,6-Me ₃ -Ph |
| 35 | 85 | Ic | Me | H | CH(OEt)cyclobutyl | 2,4,6-Me ₃ -Ph |
| | 86 | Ic | Me | H | CH(Et)cyclopropyl | 2,4,6-Me ₃ -Ph |
| | 87 | Ic | Me | H | CH(OEt)cyclopropyl | 2,4,6-Me ₃ -Ph |
| | 88 | Ic | Me | H | CH(cyclobutyl) ₂ | 2,4,6-Me ₃ -Ph |

| | | | | | | |
|----|-----|----|----|---|--|---------------------------|
| | 89 | Ic | Me | H | CH(cyclopropyl) ₂ | 2,4,6-Me ₃ -Ph |
| | 90 | Ic | Me | H | NHCH(CH ₂ OMe) ₂ | 2-Me-4-MeO-Ph |
| | 91 | Ic | Me | H | NHCH(Et)CH ₂ OMe | 2-Me-4-MeO-Ph |
| | 92 | Ic | Me | H | NH-2-butyl | 2-Me-4-MeO-Ph |
| 5 | 93 | Ic | Me | H | OCH(Et)CH ₂ OMe | 2-Me-4-MeO-Ph |
| | 94 | Ic | Me | H | O-3-pentyl | 2-Me-4-MeO-Ph |
| | 95 | Ic | Me | H | O-2-pentyl | 2-Me-4-MeO-Ph |
| | 96 | Ic | Me | H | R-2-pentyl | 2-Me-4-MeO-Ph |
| | 97 | Ic | Me | H | S-2-pentyl | 2-Me-4-MeO-Ph |
| 10 | 98 | Ic | Me | H | R-2-butyl | 2-Me-4-MeO-Ph |
| | 99 | Ic | Me | H | S-2-butyl | 2-Me-4-MeO-Ph |
| | 100 | Ic | Me | H | 3-pentyl | 2-Me-4-MeO-Ph |
| | 101 | Ic | Me | H | CH(Et)CH ₂ OH | 2-Me-4-MeO-Ph |
| | 102 | Ic | Me | H | CH(Et)CH ₂ OMe | 2-Me-4-MeO-Ph |
| 15 | 103 | Ic | Me | H | COCH ₃ | 2-Me-4-MeO-Ph |
| | 104 | Ic | Me | H | COEt | 2-Me-4-MeO-Ph |
| | 105 | Ic | Me | H | CO ₂ Et | 2-Me-4-MeO-Ph |
| | 106 | Ic | Me | H | CO-2-pentyl | 2-Me-4-MeO-Ph |
| | 107 | Ic | Me | H | CO-3-pentyl | 2-Me-4-MeO-Ph |
| 20 | 108 | Ic | Me | H | CH(OH)CH ₃ | 2-Me-4-MeO-Ph |
| | 109 | Ic | Me | H | C(OH)Me ₂ | 2-Me-4-MeO-Ph |
| | 110 | Ic | Me | H | C(OH)Ph-3-pyridyl | 2-Me-4-MeO-Ph |
| | 111 | Ic | Me | H | CH(OMe)CH ₃ | 2-Me-4-MeO-Ph |
| | 112 | Ic | Me | H | CH(OMe)Et | 2-Me-4-MeO-Ph |
| 25 | 113 | Ic | Me | H | CH(OMe)Pr | 2-Me-4-MeO-Ph |
| | 114 | Ic | Me | H | CH(OEt)CH ₃ | 2-Me-4-MeO-Ph |
| | 115 | Ic | Me | H | CH(OPr)CH ₃ | 2-Me-4-MeO-Ph |
| | 116 | Ic | Me | H | CH(OMe)Et | 2-Me-4-MeO-Ph |
| | 117 | Ic | Me | H | CH(OMe)Pr | 2-Me-4-MeO-Ph |
| 30 | 118 | Ic | Me | H | cyclobutyl | 2-Me-4-MeO-Ph |
| | 119 | Ic | Me | H | cyclopentyl | 2-Me-4-MeO-Ph |
| | 120 | Ic | Me | H | CH(Me)cyclobutyl | 2-Me-4-MeO-Ph |
| | 121 | Ic | Me | H | CH(OMe)cyclobutyl | 2-Me-4-MeO-Ph |
| | 122 | Ic | Me | H | CH(Me)cyclopropyl | 2-Me-4-MeO-Ph |
| 35 | 123 | Ic | Me | H | CH(OMe)cyclopropyl | 2-Me-4-MeO-Ph |
| | 124 | Ic | Me | H | CH(Et)cyclobutyl | 2-Me-4-MeO-Ph |
| | 125 | Ic | Me | H | CH(OEt)cyclobutyl | 2-Me-4-MeO-Ph |
| | 126 | Ic | Me | H | CH(Et)cyclopropyl | 2-Me-4-MeO-Ph |


| | | | | | | | |
|----|-----|----|----|---|--|---------------|--------|
| | 127 | Ic | Me | H | CH(OEt)cyclopropyl | 2-Me-4-MeO-Ph | |
| | 128 | Ic | Me | H | CH(cyclobutyl) ₂ | 2-Me-4-MeO-Ph | |
| | 129 | Ic | Me | H | CH(cyclopropyl) ₂ | 2-Me-4-MeO-Ph | |
| | 130 | Ic | Me | H | NHCH(CH ₂ OMe) ₂ | 2-Cl-4-MeO-Ph | |
| 5 | 131 | Ic | Me | H | NHCH(Et)CH ₂ OMe | 2-Cl-4-MeO-Ph | |
| | 132 | Ic | Me | H | NH-2-butyl | 2-Cl-4-MeO-Ph | |
| | 133 | Ic | Me | H | OCH(Et)CH ₂ OMe | 2-Cl-4-MeO-Ph | |
| | 134 | Ic | Me | H | O-3-pentyl | 2-Cl-4-MeO-Ph | |
| | 135 | Ic | Me | H | O-2-pentyl | 2-Cl-4-MeO-Ph | |
| 10 | 136 | Ic | Me | H | R-2-pentyl | 2-Cl-4-MeO-Ph | |
| | 137 | Ic | Me | H | S-2-pentyl | 2-Cl-4-MeO-Ph | |
| | 138 | Ic | Me | H | R-2-butyl | 2-Cl-4-MeO-Ph | |
| | 139 | Ic | Me | H | S-2-butyl | 2-Cl-4-MeO-Ph | |
| | 140 | Ic | Me | H | 3-pentyl | 2-Cl-4-MeO-Ph | amorph |
| 15 | 141 | Ic | Me | H | CH(Et)CH ₂ OH | 2-Cl-4-MeO-Ph | |
| | 142 | Ic | Me | H | CH(Et)CH ₂ OMe | 2-Cl-4-MeO-Ph | |
| | 143 | Ic | Me | H | COCH ₃ | 2-Cl-4-MeO-Ph | |
| | 144 | Ic | Me | H | COEt | 2-Cl-4-MeO-Ph | |
| | 145 | Ic | Me | H | CO ₂ Et | 2-Cl-4-MeO-Ph | |
| 20 | 146 | Ic | Me | H | CO-2-pentyl | 2-Cl-4-MeO-Ph | |
| | 147 | Ic | Me | H | CO-3-pentyl | 2-Cl-4-MeO-Ph | |
| | 148 | Ic | Me | H | CH(OH)CH ₃ | 2-Cl-4-MeO-Ph | |
| | 149 | Ic | Me | H | C(OH)Me ₂ | 2-Cl-4-MeO-Ph | |
| | 150 | Ic | Me | H | C(OH)Ph-3-pyridyl | 2-Cl-4-MeO-Ph | |
| 25 | 151 | Ic | Me | H | CH(OMe)CH ₃ | 2-Cl-4-MeO-Ph | |
| | 152 | Ic | Me | H | CH(OMe)Et | 2-Cl-4-MeO-Ph | |
| | 153 | Ic | Me | H | CH(OMe)Pr | 2-Cl-4-MeO-Ph | |
| | 154 | Ic | Me | H | CH(OEt)CH ₃ | 2-Cl-4-MeO-Ph | |
| | 155 | Ic | Me | H | CH(OPr)CH ₃ | 2-Cl-4-MeO-Ph | |
| 30 | 156 | Ic | Me | H | CH(OMe)Et | 2-Cl-4-MeO-Ph | |
| | 157 | Ic | Me | H | CH(OMe)Pr | 2-Cl-4-MeO-Ph | |
| | 158 | Ic | Me | H | cyclobutyl | 2-Cl-4-MeO-Ph | |
| | 159 | Ic | Me | H | cyclopentyl | 2-Cl-4-MeO-Ph | |
| | 160 | Ic | Me | H | CH(Me)cyclobutyl | 2-Cl-4-MeO-Ph | |
| 35 | 161 | Ic | Me | H | CH(OMe)cyclobutyl | 2-Cl-4-MeO-Ph | |
| | 162 | Ic | Me | H | CH(Me)cyclopropyl | 2-Cl-4-MeO-Ph | |
| | 163 | Ic | Me | H | CH(OMe)cyclopropyl | 2-Cl-4-MeO-Ph | |
| | 164 | Ic | Me | H | CH(Et)cyclobutyl | 2-Cl-4-MeO-Ph | |

| | | | | | | |
|----|-----|----|----|---|--|---------------------------|
| | 165 | Ic | Me | H | CH(OEt)cyclobutyl | 2-Cl-4-MeO-Ph |
| | 166 | Ic | Me | H | CH(Et)cyclopropyl | 2-Cl-4-MeO-Ph |
| | 167 | Ic | Me | H | CH(OEt)cyclopropyl | 2-Cl-4-MeO-Ph |
| | 168 | Ic | Me | H | CH(cyclobutyl) ₂ | 2-Cl-4-MeO-Ph |
| 5 | 169 | Ic | Me | H | CH(cyclopropyl) ₂ | 2-Cl-4-MeO-Ph |
| | 170 | Ic | Me | H | R-2-pentyl | 2-Me-6-MeO- pyrid-3-yl |
| | 171 | Ic | Me | H | S-2-pentyl | 2-Me-6-MeO- pyrid-3-yl |
| 10 | 172 | Ic | Me | H | R-2-butyl | 2-Me-6-MeO- pyrid-3-yl |
| | 173 | Ic | Me | H | S-2-butyl | 2-Me-6-MeO- pyrid-3-yl |
| | 174 | Ic | Me | H | 3-pentyl | 2-Me-6-MeO- pyrid-3-yl |
| 15 | 175 | Ic | Me | H | CH(Et)CH ₂ OH | 2-Me-6-MeO- pyrid-3-yl |
| | 176 | Ic | Me | H | CH(Et)CH ₂ OMe | 2-Me-6-MeO- pyrid-3-yl |
| 20 | 177 | Id | Me | H | NHCH(CH ₂ OMe) ₂ | 2,4-Cl ₂ -Ph |
| | 178 | Id | Me | H | NHCH(Et)CH ₂ OMe | 2,4-Cl ₂ -Ph |
| | 179 | Id | Me | H | NH-2-butyl | 2,4-Cl ₂ -Ph |
| | 180 | Id | Me | H | OCH(Et)CH ₂ OMe | 2,4-Cl ₂ -Ph |
| 25 | 181 | Id | Me | H | O-3-pentyl | 2,4-Cl ₂ -Ph |
| | 182 | Id | Me | H | O-2-pentyl | 2,4-Cl ₂ -Ph |
| | 183 | Id | Me | H | R-2-pentyl | 2,4-Cl ₂ -Ph |
| | 184 | Id | Me | H | S-2-pentyl | 2,4-Cl ₂ -Ph |
| | 185 | Id | Me | H | R-2-butyl | 2,4-Cl ₂ -Ph |
| 30 | 186 | Id | Me | H | S-2-butyl | 2,4-Cl ₂ -Ph |
| | 187 | Id | Me | H | CH(Et)CH ₂ OH | 2,4-Cl ₂ -Ph |
| | 189 | Id | Me | H | CH(Et)CH ₂ OMe | 2,4-Cl ₂ -Ph |
| | 190 | Id | Me | H | COCH ₃ | 2,4-Cl ₂ -Ph |
| | 191 | Id | Me | H | COEt | 2,4-Cl ₂ -Ph |
| 35 | 192 | Id | Me | H | CO ₂ Et | 2,4-Cl ₂ -Ph |
| | 193 | Id | Me | H | CO-2-pentyl | 2,4-Cl ₂ -Ph |
| | 194 | Id | Me | H | CO-3-pentyl | 2,4-Cl ₂ -Ph |
| | 195 | Id | Me | H | CH(OH)CH ₃ | 2,4-Cl ₂ -Ph |

| | | | | | | | |
|----|-----|----|----|---|--|-------------------------------|-----------|
| | 196 | Id | Me | H | C(OH)Me ₂ | 2,4-Cl ₂ -Ph | |
| | 197 | Id | Me | H | C(OH)Ph-3-pyridyl | 2,4-Cl ₂ -Ph | |
| | 198 | Id | Me | H | CH(OMe)CH ₃ | 2,4-Cl ₂ -Ph | |
| | 199 | Id | Me | H | CH(OMe)Et | 2,4-Cl ₂ -Ph | |
| 5 | 200 | Id | Me | H | CH(OMe)Pr | 2,4-Cl ₂ -Ph | |
| | 201 | Id | Me | H | CH(OEt)CH ₃ | 2,4-Cl ₂ -Ph | |
| | 202 | Id | Me | H | CH(OPr)CH ₃ | 2,4-Cl ₂ -Ph | |
| | 203 | Id | Me | H | CH(OMe)Et | 2,4-Cl ₂ -Ph | |
| | 204 | Id | Me | H | CH(OMe)Pr | 2,4-Cl ₂ -Ph | |
| 10 | 205 | Id | Me | H | cyclobutyl | 2,4-Cl ₂ -Ph | |
| | 206 | Id | Me | H | cyclopentyl | 2,4-Cl ₂ -Ph | |
| | 207 | Id | Me | H | CH(Me)cyclobutyl | 2,4-Cl ₂ -Ph | |
| | 208 | Id | Me | H | CH(OMe)cyclobutyl | 2,4-Cl ₂ -Ph | |
| | 209 | Id | Me | H | CH(Me)cyclopropyl | 2,4-Cl ₂ -Ph | |
| 15 | 210 | Id | Me | H | CH(OMe)cyclopropyl | 2,4-Cl ₂ -Ph | |
| | 211 | Id | Me | H | CH(Et)cyclobutyl | 2,4-Cl ₂ -Ph | |
| | 212 | Id | Me | H | CH(OEt)cyclobutyl | 2,4-Cl ₂ -Ph | |
| | 213 | Id | Me | H | CH(Et)cyclopropyl | 2,4-Cl ₂ -Ph | |
| | 214 | Id | Me | H | CH(OEt)cyclopropyl | 2,4-Cl ₂ -Ph | |
| 20 | 215 | Id | Me | H | CH(cyclobutyl) ₂ | 2,4-Cl ₂ -Ph | |
| | 216 | Id | Me | H | CH(cyclopropyl) ₂ | 2,4-Cl ₂ -Ph | |
| | 217 | Ic | Me | H | 3-pentyl | 2,4-Me ₂ -4-MeOPh | 80-82 |
| | 218 | Ic | Me | - | 4-heptyl | 2,4-Cl ₂ -Ph | 75.5-76.5 |
| | 219 | Ic | Me | - | CH(Me)cyclobutyl | 2,4-Cl ₂ -Ph | |
| 25 | 220 | Ic | Me | - | CH(CH ₂ OMe)Pr | 2,4-Cl ₂ -Ph | |
| | 221 | Ic | Me | - | CH(CH ₂ CH ₂ OMe) ₂ | 2,4-Cl ₂ -Ph | |
| | 222 | Ic | Me | - | CH(CH ₂ CH ₂ OMe)Et | 2,4-Cl ₂ -Ph | |
| | 223 | Ic | Me | - | CH(CH ₂ CH ₂ OMe)- cyclobutyl | 2,4-Cl ₂ -Ph | |
| 30 | 224 | Ic | Me | - | CH(CH ₂ CH ₂ OMe) 3-tetrahydrofuranyl | 2,4-Cl ₂ -Ph | |
| | 225 | Ic | Me | - | 3-pentyl | 2-Cl-4-OMe-Ph | |
| | 226 | Ic | Me | - | CH(Me)Pr | 2-Cl-4-OMe-Ph | |
| | 227 | Ic | Me | - | 4-heptyl | 2-Cl-4-OMe-Ph | |
| 35 | 228 | Ic | Me | - | CH(CH ₂ OMe)Pr | 2-Cl-4-OMe-Ph | |
| | 229 | Ic | Me | - | CH(Me)cyclobutyl | 2-Cl-4-OMe-Ph | |
| | 230 | Ic | Me | - | CH(Me)Pr | 2-Cl-4,5-OMe ₂ -Ph | |
| | 231 | Ic | Me | - | CH(Me)cyclobutyl | 2-Cl-4,5-OMe ₂ -Ph | |

| | | | | | | | |
|----|-----|----|----|---|--|-------------------------------|-----------|
| | 232 | Ic | Me | - | CH(CH ₂ CH ₂ OMe) ₂ | 2-Cl-4,5-OMe ₂ -Ph | |
| | 233 | Ic | Me | - | 4-heptyl | 2-Cl-4,5-OMe ₂ -Ph | |
| | 234 | Ic | Me | - | CH(CH ₂ OMe) Pr | 2-Cl-4,5-OMe ₂ -Ph | |
| | 235 | Ic | Me | - | CH(CH ₂ CH ₂ OMe) | 2-Cl-4,5-OMe ₂ -Ph | |
| 5 | | | | | cyclobutyl | | |
| | 236 | Ic | Me | - | CH(CH ₂ CH ₂ OMe) - | 2-Cl-4,5-OMe ₂ -Ph | |
| | | | | | 3-tetrahydrofuranlyl | | |
| | 237 | Ic | Me | - | CH(Me) cyclobutyl | 2-Cl-4-OMe-5-FPh | |
| | 238 | Ic | Me | - | CH(Me) Pr | 2-Cl-4-OMe-5-F-Ph | |
| 10 | 239 | Ic | Me | - | CH(CH ₂ CH ₂ OMe) ₂ | 2-Cl-4-OMe-5-F-Ph | |
| | 240 | Ic | Me | - | CH(CH ₂ CH ₂ OMe) Et | 2-Cl-4-OMe-5-F-Ph | |
| | 241 | Ic | Me | - | CH(CH ₂ OMe) Pr | 2-Cl-4-OMe-5-F-Ph | |
| | 242 | Ic | Me | - | CH(CH ₂ CH ₂ OMe) - | 2-Cl-4-OMe-5-F-Ph | |
| | | | | | cyclobutyl | | |
| 15 | 243 | Ic | Me | - | CH(CH ₂ CH ₂ OMe) - | 2-Cl-4-OMe-5-F-Ph | |
| | | | | | 3-tetrahydrofuranlyl | | |
| | 244 | Ic | Me | - | 3-pentyl | 2-Cl-4-OEt-Ph | |
| | 245 | Ic | Me | - | CH(Me) Pr | 2-Cl-4-OEt-Ph | |
| | 246 | Ic | Me | - | 4-heptyl | 2-Cl-4-OEt-Ph | 85.5-86.5 |
| 20 | 247 | Ic | Me | - | CH(Me) cyclobutyl | 2,4-OMe ₂ -Ph | |
| | 248 | Ic | Me | - | 3-pentyl | 2,4-OMe ₂ -Ph | |
| | 249 | Ic | Me | - | 4-heptyl | 2,4-OMe ₂ -Ph | 87-88 |
| | 250 | Ic | Me | - | 3-pentyl | 2-Me-4-Cl-Ph | |
| | 251 | Ic | Me | - | 3-pentyl | 4-OMe-Ph | |
| 25 | 252 | Ic | Me | - | 3-pentyl | 4-Cl-Ph | |
| | 253 | Ic | Me | - | 3-pentyl | 2,5-Me ₂ -4-OMe-Ph | 80-82 |
| | 254 | Ic | Me | - | 3-pentyl | 2-Cl-4-SO ₂ Me-Ph | |
| | 255 | Ic | Me | - | 3-pentyl | 2-Me-4-NMe ₂ -Ph | |
| | 256 | Ic | Me | - | 3-pentyl | 2-Cl-4-NMe ₂ -Ph | |
| 30 | 257 | Ic | Me | - | 3-pentyl | 2-CF ₃ -4-F-Ph | |
| | 258 | Ic | Me | - | 3-pentyl | 2-OMe-4-Me-Ph | |
| | 259 | Ic | Me | - | CH(Me) Pr | 2-OH-4-OMe-Ph | |
| | 260 | Ic | Me | - | CH(Me) Pr | 2-Me-4-OMe-5-F-Ph | |
| | 261 | Id | Me | - | 3-pentyl | 2,4-Cl ₂ -Ph | |
| 35 | 262 | Id | Me | - | 3-pentyl | 2-Cl-4,5-OMe ₂ -Ph | |

TABLE 1a



| 5 | Ex. | Formula | R^5 | R^4 | R^2 | R^1 | mp (°C) |
|---|-----|---------|-------|-------|----------|-------------------------------|---------|
| | 263 | | Me | - | 3-pentyl | 2-Cl-4,5-OMe ₂ -Ph | 249-250 |

10

In addition to examples 1-10, additional examples 11-248 and additional examples having the following R^4 (or R^5), R^2 and R^3 were or may readily be prepared according to the procedures described herein.

The preferred groups for R^5 in the compounds of formula Ic and Id are methyl (Me). For R^2 , in compounds of formula Ic and Id having R^5 as Me (or C₂-C₆alkyl), the preferred groups include 3-pentyl, NEt₂, butyl, NHCH(CH₂OMe)₂, NHCH(CH₂OEt)₂, NHCH(Et)CH₂OMe, NH-3-heptyl, NH-3-pentyl, NH-2-butyl, NH-3-hexyl, NHCH(CH₂Ph)CH₂OMe, NHCH(Et)CH₂CH₂OMe, NH-cyclobutyl, NH-cyclopentyl, NEtPr, NEtBu, NMePr, NMePh, NPr₂, NPr(CH₂-c-C₃H₅), N(CH₂CH₂OMe)₂, morpholino, N(CH₂Ph)CH₂CH₂OMe, N(Me)CH₂CH₂OMe, N(Et)CH₂CH₂OMe, N(CH₂-c-C₃H₅)CH₂CH₂OMe, N(CH₂-c-C₃H₅)Pr, N(CH₂-c-C₃H₅)Et, OEt, OCH(Et)CH₂OMe, OCH(Et)CH₂CH₂OMe, OCH(Me)CH₂CH₂OMe, O-3-pentyl, O-2-pentyl, S-3-pentyl, S-2-pentyl, SEt, S(O)Et, SO₂Et, S-3-pentyl, S(O)-3-pentyl, SO₂-3-pentyl, S-2-pentyl, S(O)-2-pentyl, SO₂-2-pentyl, CH(CO₂Et)₂, C(Et)(CO₂Et)₂, CH(Et)CH₂OH, CH(Et)CH₂OMe, CH(Et)CH₂CH₂OMe, CONMe₂, COCH₃, COEt, COPr, CO-2-pentyl, CO-3-pentyl, CH(OH)CH₃, C(OH)Me₂, C(OH)Ph-3-pyridyl, CH(OMe)CH₃, CH(OMe)Et, CH(OMe)Pr, CH(OEt)CH₃, CH(OPr)CH₃, 2-pentyl, 2-butyl, cyclobutyl, cyclopentyl, CH(Me)cyclobutyl, CH(OMe)cyclobutyl, CH(OH)cyclobutyl, CH(Me)cyclopropyl,

- CH(OMe)cyclopropyl, CH(OH)cyclopropyl, CH(Et)cyclobutyl, CH(Et)cyclopropyl, CH(OMe)cyclobutyl, CH(OMe)cyclopropyl, CH(OEt)cyclobutyl, CH(OEt)cyclopropyl, CH(Me)CH₂-cyclobutyl, CH(OMe)CH₂-cyclobutyl, CH(OH)CH₂-cyclobutyl,
- 5 CH(Me)CH₂-cyclopropyl, CH(OMe)CH₂-cyclopropyl, CH(OH)CH₂-cyclopropyl, CH(Et)CH₂-cyclobutyl, CH(Et)CH₂-cyclopropyl, CH(OMe)CH₂-cyclobutyl, CH(OMe)CH₂-cyclopropyl, CH(OEt)CH₂-cyclobutyl, CH(OEt)CH₂-cyclopropyl, CH(CH₂OMe)cyclobutyl, CH(CH₂OMe)cyclopropyl,
- 10 CH(CH₂OEt)cyclobutyl, CH(CH₂OEt)cyclopropyl, CH(cyclobutyl)₂, CH(cyclopropyl)₂, CH(Et)CH₂CONMe₂, CH(Et)CH₂CH₂NMe₂, CH(CH₂OMe)Me, CH(CH₂OMe)Et, CH(CH₂OMe)Pr, CH(CH₂OEt)Me, CH(CH₂OEt)Et, CH(CH₂OEt)Pr, CH(CH₂C≡CMe)Et, CH(CH₂C≡CMe)Et. The preferred groups for
- 15 R³ with R² and R⁵ as defined above include 2,4-Cl₂-Ph, 2,4,6-Me₃-Ph, 2,4-Me₂-Ph, 2-Me-4-MeO-Ph, 2-Cl-4-MeO-Ph, 2-Cl-4,5-(MeO)₂-Ph, 2-Cl-4-MeO-5-F-Ph, 2-Me-4-MeO-5-F-Ph, 2,5-(Me)₂-4-MeO-Ph, 2-Me-4-NMe₂-Ph, 2-CF₃-4-MeO-Ph, 2-Me-4-(COMe)-Ph, 2-Me-6-Me₂N-pyrid-3-yl, 4-Me-2-Me₂N-pyrid-5-
- 20 yl, 2-Me-6-MeO-pyrid-3-yl, 4-Me-2-MeO-pyrid-5-yl. Each of the compounds within the independent generic variations may readily be prepared according to the procedures described in Schemes 1-3 and 5.
- For compounds of formulas Ie and If, in addition to
- 25 example 10 in Table 1, the above variables for Ic and Id may be used in the compounds of formulas Ie and If except that R⁵ groups are used as the R⁴ variables. The following compounds may be prepared as the preferred embodiments wherein R⁴ is selected from H, OCH₃, CH₃, and
- 30 C₂H₅; R² is selected from CH(C₂H₅)₂, CH(c-C₃H₅)₂, CHC₂H₅(c-C₃H₅), CH(C₂H₅)₂, CH(c-C₃H₅)₂; and R³ is selected from 2,4-Cl₂-Ph, 2-Cl-4-CH₃O-Ph, 2,4,6-(CH₃)₃-Ph, 2-Cl-4-CF₃-Ph and 2-(CH₃)₂N-4-CH₃-pyridin-5-yl. In addition, the methoxy group or the ethyl group in the R¹ position also
- 35 preferably includes CH₃ and H. Each of the compounds within the independent generic variations may readily be prepared according to the procedure described in Scheme 4. Compounds of formula IIa and IIb are readily prepared

according to the procedure described in Scheme 6. These compounds also preferably have the variables shown in the examples and described above.

5

Utility

CRF-R1 Receptor Binding Assay for the Evaluation of Biological Activity

10

Radioligand binding experiments

Compounds of the invention were tested for in vitro activity as CRF receptor antagonists. The tests described below demonstrated that the examples tested had K_i s of 10,000 nM or less and are thus useful as CRF receptor antagonists. Preferred antagonists have or will have a K_i of 1,000 nM or less. Radioligand binding experiments were performed with membranes from rat frontal cortex to determine binding affinities (K_i 's) of test compounds for the rat CRH₁ receptor using a modified version of methods described earlier (see E.B. DeSouza, J. Neurosci, 7:88, 1987). Rat cortex was homogenized in tissue buffer (containing 50 mM HEPES, 10 mM MgCl₂, 2 mM EGTA, and 1 µg/ml each of aprotinin, leupeptin, and pepstatin, pH 7.0 @ 23°C) using a Brinkman Polytron (PT-10, setting 6 for 10 sec). The homogenate was centrifuged at 48,000 X g for 12 min and the resulting pellet was washed by two sequential re-suspension and centrifugation steps. The final pellet was suspended to tissue buffer to a working concentration of 0.1 mg/ml protein. Protein determinations were made using the bicinchoninic acid (BCA) assay (Pierce, Rockford, IL) with bovine serum albumin as the standard.

All test compounds were prepared in assay buffer, which was identical to the tissue buffer except for the inclusion of 0.15 mM bacitracin and 0.1% w/v ovalbumin. Binding assay were conducted in disposable polypropylene 96-well plates (Costar Corp., Cambridge, MA) and initiated by the addition of 100 µl membrane

homogenate (containing 40-60 µg protein) to 200 µl of assay buffer containing radioligands (150 pM, final concentration, [¹²⁵I] tyr^o ovine CRH; New England Nuclear, MA) and competing test compounds. Specific
5 binding was determined in the presence of 10 µM α-helical CRH. Competition experiments were conducted using 12 concentrations of ligand (ranging from 1 X 10⁻¹¹ to 1 X 10⁻⁵ M). The reactions mixtures were incubated to equilibrium for 2 hr at 23°C and terminated by rapid
10 filtration using a cell harvester (Inotech Biosystems Inc., Lansing MI) over GFF glass-fibers (pre-soaked in 0.3 % v/v polyethyleneimine). Filters were rapidly washed 3X with 0.3 ml cold wash buffer (PBS, pH 7.0, containing 0.01% Triton X-100), dried, and counted in a
15 gamma counter at 80% efficiency.

Binding affinities (K_i's) of ligands for the CRH₁ receptor were calculated using the iterative nonlinear regression curve-fitting programs (LIGAND) of Munson and Rodbard (Anal. Biochem. 1980, 107, 220-239) or
20 Prism (GraphPad Prism, San Diego, CA). Data were best-fit by the one-site/state competition equation.

Inhibition of CRF-Stimulated Adenylate Cyclase Activity

Inhibition of CRF-stimulated adenylate cyclase
25 activity can be performed as described by G. Battaglia et al. *Synapse* 1:572 (1987). Briefly, assays are carried out at 37° C for 10 min in 200 µl of buffer containing 100 mM Tris-HCl (pH 7.4 at 37° C), 10 mM MgCl₂, 0.4 mM EGTA, 0.1% BSA, 1 mM
30 isobutylmethylxanthine (IBMX), 250 units/ml phosphocreatine kinase, 5 mM creatine phosphate, 100 mM guanosine 5'-triphosphate, 100 nM oCRF, antagonist peptides (concentration range 10⁻⁹ to 10⁻⁶M) and 0.8 mg original wet weight tissue (approximately 40-60 mg
35 protein). Reactions are initiated by the addition of 1 mM ATP/³²P]ATP (approximately 2-4 mCi/tube) and terminated by the addition of 100 µl of 50 mM Tris-HCl, 45 mM ATP and 2% sodium dodecyl sulfate. In order to

monitor the recovery of cAMP, 1 μ l of [3 H]cAMP (approximately 40,000 dpm) is added to each tube prior to separation. The separation of [32 P]cAMP from [32 P]ATP is performed by sequential elution over Dowex and alumina columns.

In vivo Biological Assay

The *in vivo* activity of the compounds of the present invention can be assessed using any one of the biological assays available and accepted within the art. Illustrative of these tests include the Acoustic Startle Assay, the Stair Climbing Test, and the Chronic Administration Assay. These and other models useful for the testing of compounds of the present invention have been outlined in C.W. Berridge and A.J. Dunn *Brain Research Reviews* 15:71 (1990).

Compounds may be tested in any species of rodent or small mammal.

20

Compounds of this invention have utility in the treatment of imbalances associated with abnormal levels of corticotropin releasing factor in patients suffering from depression, affective disorders, and/or anxiety.

25

Compounds of this invention can be administered to treat these abnormalities by means that produce contact of the active agent with the agent's site of action in the body of a mammal. The compounds can be administered by any conventional means available for use in conjunction with pharmaceuticals either as individual therapeutic agent or in combination of therapeutic agents. They can be administered alone, but will generally be administered with a pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

30

35

The dosage administered will vary depending on the use and known factors such as pharmacodynamic character of the particular agent, and its mode and route of

administration; the recipient's age, weight, and health; nature and extent of symptoms; kind of concurrent treatment; frequency of treatment; and desired effect. For use in the treatment of said
5 diseases or conditions, the compounds of this invention can be orally administered daily at a dosage of the active ingredient of 0.002 to 200 mg/kg of body weight. Ordinarily, a dose of 0.01 to 10 mg/kg in divided doses one to four times a day, or in sustained release
10 formulation will be effective in obtaining the desired pharmacological effect.

Dosage forms (compositions) suitable for administration contain from about 1 mg to about 100 mg of active ingredient per unit. In these pharmaceutical
15 compositions, the active ingredient will ordinarily be present in an amount of about 0.5 to 95% by weight based on the total weight of the composition.

The active ingredient can be administered orally is solid dosage forms, such as capsules, tablets and
20 powders; or in liquid forms such as elixirs, syrups, and/or suspensions. The compounds of this invention can also be administered parenterally in sterile liquid dose formulations.

Gelatin capsules can be used to contain the active
25 ingredient and a suitable carrier such as but not limited to lactose, starch, magnesium stearate, steric acid, or cellulose derivatives. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release
30 products to provide for continuous release of medication over a period of time. Compressed tablets can be sugar-coated or film-coated to mask any unpleasant taste, or used to protect the active ingredients from the atmosphere, or to allow selective
35 disintegration of the tablet in the gastrointestinal tract.

Liquid dose forms for oral administration can contain coloring or flavoring agents to increase patient acceptance.

In general, water, pharmaceutically acceptable
5 oils, saline, aqueous dextrose (glucose), and related
sugar solutions and glycols, such as propylene glycol
or polyethylene glycol, are suitable carriers for
parenteral solutions. Solutions for parenteral
administration preferably contain a water soluble salt
10 of the active ingredient, suitable stabilizing agents,
and if necessary, buffer substances. Antioxidizing
agents, such as sodium bisulfite, sodium sulfite, or
ascorbic acid, either alone or in combination, are
suitable stabilizing agents. Also used are citric acid
15 and its salts, and EDTA. In addition, parenteral
solutions can contain preservatives such as
benzalkonium chloride, methyl- or propyl-paraben, and
chlorobutanol.

Suitable pharmaceutical carriers are described in
20 "Remington's Pharmaceutical Sciences", A. Osol, a
standard reference in the field.

Useful pharmaceutical dosage-forms for
administration of the compounds of this invention can
be illustrated as follows:

25

Capsules

A large number of units capsules are prepared by
filling standard two-piece hard gelatin capsules each
with 100 mg of powdered active ingredient, 150 mg
30 lactose, 50 mg cellulose, and 6 mg magnesium stearate.

Soft Gelatin Capsules

A mixture of active ingredient in a digestible oil
such as soybean, cottonseed oil, or olive oil is
35 prepared and injected by means of a positive
displacement was pumped into gelatin to form soft
gelatin capsules containing 100 mg of the active
ingredient. The capsules were washed and dried.


Tablets

- A large number of tablets are prepared by conventional procedures so that the dosage unit was 100 mg active ingredient, 0.2 mg of colloidal silicon
- 5 dioxide, 5 mg of magnesium stearate, 275 mg of microcrystalline cellulose, 11 mg of starch, and 98.8 mg lactose. Appropriate coatings may be applied to increase palatability or delayed adsorption.
- 10 The compounds of this invention may also be used as reagents or standards in the biochemical study of neurological function, dysfunction, and disease. The preferred indication and use for the compounds and compositions of the invention is in the treatment of
- 15 depression or anxiety.

WHAT IS CLAIMED IS:

1. A compound of formula I:

5



or a stereoisomer or pharmaceutically acceptable salt thereof, wherein:

10

A equals N or CR⁵;

B equals N or CR⁴,

provided that both A and B can not be N or provided that

15

A can not be CR⁵ and B can not be CR⁴ to form a pyrazolopyrimidine;

R¹ is independently selected from the group consisting of

20

H,
halogen,

CN,

C₁₋₆ alkyl,

C₂₋₁₀ alkenyl,

25

C₂₋₁₀ alkynyl,

C₃₋₆ cycloalkyl,

C₁₋₆ alkyloxy,

C₁₋₆ alkylS(O)_n,

-NR^{1a}R^{1b} wherein R^{1a} and R^{1b} are independently selected from

30

H, C₁₋₄ alkyl, C₃₋₆ cycloalkyl, -C(O)C₁₋₄alkyl,

C₁₋₆ alkylNR^{1a}R^{1b},

NR^{1a}COR^{1b},

-C(O)NR^{1a}R^{1b},

-O-C(O)C₁₋₄alkyl,

-XR^{1c} wherein R^{1c} is selected from H or -C₁₋₄ alkylaryl;

5 X is selected from O or S(O)_n,

wherein R¹ is substituted with 0-6 substituents selected from halogen, C₁₋₄ alkyl, C₃₋₈ cycloalkyl, C₁₋₆ alkyloxy, C₁₋₄ haloalkyl, C₁₋₄ alkylamino, C₂₋₈ dialkylamino, C₁₋₄ alkylthio,
10 C₁₋₄ alkylsulfinyl or C₁₋₄ alkylsulfonyl;

R² is selected from the group consisting of
H, OR⁷, SH, NR⁶R⁷, C(OH)R⁶R^{6a}, C(OR⁷)R⁶R^{6a}, S(O)_nR¹³, COR⁷,
CO₂R⁷, CHR⁶(OR⁷)R^{6a}, OC(O)R¹³, NO, NO₂, NR⁶C(O)R⁷, N(COR⁷)₂,
15 NR⁸CONR⁶R⁷, NR⁶CO₂R⁷; or

C₁₋₁₀ alkyl,

C₂₋₁₀ alkenyl,

C₂₋₁₀ alkynyl,

20 C₃₋₈ cycloalkyl,

C₃₋₆ cycloalkyl C₁₋₆ alkyl,

C₁₋₁₀ alkyloxy,

C₁₋₁₀ alkyloxyC₁₋₁₀ alkyl,

-SO₂-C₁₋₁₀alkyl

25 -SO₂R^{2a} wherein R^{2a} is aryl,

-SO₂R^{2b} wherein R^{2b} is heteroaryl,

-NR^{2c}R^{2d} wherein R^{2c} and R^{2d} are independently selected from
H, C₁₋₈ alkyl, S(O)_nC₁₋₄alkyl, C(O)NR^{2c}R^{2d}, CO₂C₁₋₄alkyl,
C₃₋₈ cycloalkyl, C₁₋₆ alkyloxyC₁₋₆ alkyl, -C(O)C₁₋₄alkyl
30 or R^{2c} and R^{2d} may join to form a heterocyclic ring
having 0-3 heteroatoms selected from O, N or S,

- halogen,

-CN,

35 -C(O)-L wherein L is selected from H, NR^{2c}R^{2d}, C₁₋₆ alkyl or
OC₁₋₄ alkyl, O(CH₂)_mOR wherein R is C₁₋₃ alkyl,
O(CH₂)_m-NR^{2c}R^{2d}, OH, C(O)OC₁₋₆alkyl or aryl or heteroaryl
wherein m is 1-4;

-OC(O)-M wherein M is selected from C₁₋₄ alkyl, C₁₋₄
haloalkyl, C₂₋₈ alkoxyalkyl, C₃₋₆ cycloalkyl, C₄₋₁₂
cycloalkylalkyl, aryl, C₁₋₆ alkylaryl, heteroaryl, C₁₋₆
5 alkylheteroaryl;

n is 0, 1 or 2; and wherein

R² is substituted with 0-3 substituents independently
10 selected from R', R'', R''' wherein R', R'' and R''' are
independently selected from C₁₋₆ alkyl, C₃₋₇ cycloalkyl,
hydroxyc₁₋₆ alkyl, C₁₋₆ alkyloxy C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆
alkynyl, C₁₋₆ alkyloxy, hydroxy, or

15 R² is substituted with 0-3 substituents independently
selected from:

halogen,

-CN,

20 -S(O)_nR^{2o} wherein R^{2o} is selected from C₁₋₄ alkyl, C₁₋₄
haloalkyl, C₁₋₄ alkyloxy C₁₋₄ alkyl, C₃₋₆ cycloalkyl;

-COR^{2f} wherein R^{2f} is selected from H, C₁₋₄ alkyl, C₁₋₄
haloalkyl, C₁₋₄ alkyloxy C₁₋₄ alkyl, C₃₋₆
25 cycloalkyl, and C₃₋₆ cycloalkyl C₁₋₄ alkyl;

-CO₂R^{2f},

-NR^{2g}COR^{2f} wherein R^{2g} is selected from H, C₁₋₆ alkyl, C₃₋₇
cycloalkyl, C₃₋₆ cycloalkyl C₁₋₆ alkyl;

30 -N(COR^{2f})₂,

-NR^{2g}CONR^{2f}R^{2h}, wherein R^{2h} is selected from H, C₁₋₆ alkyl,
C₁₋₄ haloalkyl, C₁₋₄ alkoxy C₁₋₄ alkyl,
C₃₋₆ cycloalkyl and C₃₋₆ cycloalkyl C₁₋₆
alkyl;

35

-NR^{2g}CO₂R^{2o},

-CONR^{2g}R^{2h},

1-morpholinyl,

- 1-piperidinyl,
1-piperazinyl,
and
C₃₋₈ cycloalkyl wherein 0-1 carbon atoms in the C₄₋₈
5 cycloalkyl is replaced by a group selected from
-O-, -S(O)_n-, -NR^{2g}-, -NCO₂R^{2o}, -NCOR^{2o},
and -NSO₂R^{2o}; and wherein N₄ in
1-piperazinyl is substituted with 0-1
substituents selected from R^{2g}, CO₂R^{2o}, COR^{2o} and
10 SO₂R^{2o}; or
- the group R²ⁱ, R^{2j}, R^{2k}, C₁₋₆ alkyl, C₂₋₈
alkenyl, C₂₋₈ alkynyl, Br, Cl, F, I, C₁₋₄ haloalkyl, -OR^{2g},
-NR^{2g}R^{2h}, -C₁₋₆ alkyl-OR^{2g}, and C₃₋₈ cycloalkyl which is
15 substituted with 0-1 R²ⁱ and in which 0-1 carbons of C₄₋₈
cycloalkyl is replaced by -O-, wherein
R²ⁱ is selected from aryl wherein aryl includes
phenyl, naphthyl, indanyl and indenyl, each
R²ⁱ being substituted with 0-1 OR^{2m} and 0-5
20 substituents independently selected from the group C₁₋₆
alkyl, C₃₋₆ cycloalkyl, Br, Cl, F, I, C₁₋₄ haloalkyl, -CN,
nitro, -SH, -S(O)_nR²ⁿ, -COR^{2m}, -OC(O)R²ⁿ, -NR^{2g}COR^{2m}, -
N(COR^{2m})₂,
-NR^{2g}CONR^{2o}R^{2p}, -NR^{2g}CO₂R²ⁿ, -NR^{2o}R^{2p} and -CONR^{2o}R^{2p};
25
- R^{2j} is selected from heteroaryl wherein heteroaryl
includes pyridyl, pyrimidinyl, triazinyl, furanyl,
quinolinyl, isoquinolinyl, thienyl, imidazolyl,
thiazolyl, indolyl, pyrrolyl, oxazolyl, benzofuranyl,
30 benzothienyl, benzothiazolyl, benzoxazolyl, isoxazolyl,
pyrazolyl, triazolyl, tetrazolyl, indazolyl, 2,3-
dihydrobenzofuranyl, 2,3-dihydrobenzothienyl, 2,3-
dihydrobenzothienyl-s-oxide, 2,3-dihydro-benzothienyl-s-
dioxide, indolinyl, benzoxazolin-2-onyl, benzodioxolanyl
35 and benzodioxane, each heteroaryl being substituted on 0-
4 carbon atoms with a substituent independently selected
from the group C₁₋₆ alkyl, C₃₋₆ cycloalkyl, Br, Cl, F, I, C₁₋₄
haloalkyl, -CN, nitro, OR^{2m}, -SH, -S(O)_nR^{2h}, -COR²ⁿ, -

OC(O)R^{2h}, -NR^{2g}COR^{2m}, -N(COR^{2m})₂, -NR^{2g}CONR^{2o}R^{2p}, -NR^{2g}CO₂R^{2h}, -NR^{2o}R^{2p} and -CONR^{2o}R^{2p} and each heteroaryl being substituted on any nitrogen atom with 0-1 substituents selected from the group R^{2g}, CO₂R^{2o}, COR^{2o} and SO₂R^{2o};

5

R^{2k} is heterocyclyl which is a saturated or partially saturated heteroaryl as defined for R^{2j}, each heterocyclyl being substituted on 0-4 carbon atoms with a substituent independently selected from the group C₁₋₆ alkyl, C₃₋₆ cycloalkyl, Br, Cl, F, I, C₁₋₄ haloalkyl, -CN, nitro, -OR^{2m}, -SH, -S(O)_nR^{2h}, -COR^{2m}, -OC(O)R^{2h}, -NR^{2g}COR^{2m}, -N(COR^{2m})₂, -NR^{2g}CONR^{2o}R^{2p}, NR^{2g}CO₂R^{2h}, -NR^{2o}R^{2p} and -CONR^{2o}R^{2p} and each heterocyclyl being substituted on any nitrogen atom with 0-1 substituents selected from the group R^{2f}, CO₂R^{2o}, COR^{2o} and SO₂R^{2o};

15

wherein

R²¹ is H, C₁₋₄ alkyl, C₃₋₆ cycloalkyl-C₁₋₄ alkyl and C₃₋₈ cycloalkyl;

20

R^{2m} is H, C₁₋₆ alkyl, C₃₋₆ cycloalkyl C₁₋₆ alkyl, C₁₋₂ alkyloxy C₁₋₂ alkyl, C₁₋₄ haloalkyl, R^{2g}S(O)_n-C₁₋₄ alkyl and R^{2r}R^{2s}N-C₂₋₄ alkyl;

25

R²ⁿ is H, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, C₃₋₆ cycloalkyl-C₁₋₆ alkyl, C₁₋₂ alkyloxy C₁₋₂ alkyl, and C₁₋₄ haloalkyl;

R^{2o} and R^{2p} are independently selected at each occurrence from H, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, C₃₋₆ cycloalkyl C₁₋₆ alkyl and C₁₋₄ haloalkyl;

30

R^{2g} is selected from C₁₋₆ alkyl, C₁₋₄ haloalkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, C₃₋₆ cycloalkyl, C₃₋₆ cycloalkyl-C₁₋₆ alkyl, aryl, aryl(C₁₋₄ alkyl), heteroaryl and heteroaryl (C₁₋₄ alkyl)- and benzyl, each benzyl being substituted on the aryl moiety with 0-1 substituents selected from the group C₁₋₄

35

alkyl, Br, Cl, F, I, C₁₋₄ haloalkyl, nitro, C₁₋₄ alkoxy C₁₋₄ haloalkoxy, and dimethylamino;

5 R^{2r}R^{2s} taken together with the N form 1-pyrrolidinyl, 1-morpholinyl, 1-piperidinyl or 1-piperazinyl wherein N₄ in 1-piperiazinyl is substituted with 0-1 substituents selected from the group R^{2t}, CO₂R^{2q}, COR^{2q} and SO₂R^{2q};

10 R^{2t} is selected from H, C₁₋₄ alkyl, C₁₋₄ haloalkyl, C₁₋₄ alkoxy -C₁₋₄ alkyl, C₃₋₆ cycloalkyl, C₃₋₆ cycloalkyl - C₁₋₆ alkyl, aryl, aryl (C₁₋₄ alkyl)-, heteroaryl and heteroaryl (C₁₋₄ alkyl);

15 R³ is selected from an aryl or heteroaryl group attached through an unsaturated carbon atom;

aryl is selected from phenyl, naphthyl, indanyl and indenyl, each aryl being substituted with 0-5 substituents independently selected at each occurrence
20 from C₁₋₆ alkyl, C₃₋₆ cycloalkyl, methylenedioxy, C₁₋₄ alkyloxy-C₁₋₄ alkyloxy, -OR^{2m}, Br, Cl, F, I, C₁₋₄ haloalkyl, -CN, -NO₂, -SH, -S(O)_nR²ⁿ, -COR^{2m}, -CO₂R^{2m}, -OC(O)R²ⁿ, -NR^{2g}COR^{2m}, -N(COR^{2m})₂, -NR^{2g}CONR^{2o}R^{2p}, -NR^{2g}CO₂R^{2h}, -NR^{2o}R^{2p} and CONR^{2o}R^{2p};

25 heteroaryl is selected from the group pyridyl, pyrimidyl, triazinyl, furanyl, quinolinyl, isoquinolinyl, thienyl, imidazolyl, thiazolyl, indolyl, pyrrolyl, oxazolyl, benzofuranyl, benzothienyl, benzothiazolyl, benzoxazolyl,
30 isoxazolyl, triazolyl, tetrazolyl, indazolyl, 2,3-dihydrobenzo-furanyl, 2,3-dihydrobenzothienyl, 2,3-dihydro-benzothienyl-S-oxide, 2,3-dihydrobenzothienyl-s-dioxide, indolinyl, benzoxazolin-2-on-yl, benzodioxolanyl and benzodioxane, each heteroaryl being substituted at 0-
35 4 carbon atoms with a substituent independently selected at each occurrence from the group C₁₋₆ alkyl, C₃₋₆ cycloalkyl, Br, F, I, C₁₋₄ haloalkyl, -CN, NR^{2g}R^{2h}, nitro, -OR^{2m}, -SH, -S(O)_nR²ⁿ, COR^{2m}, -CO₂R^{2m}, -OC(O)R²ⁿ, -NR^{2g}COR^{2m}, -

$N(COR^{2m})_2$, $-NR^{2g}CONR^{2o}R^{2p}$ and each heteroaryl being substituted at any nitrogen atom with 0-1 substituents selected from the group R^{2g} , CO_2R^{3a} , COR^{3a} and SO_2R^{3a} wherein,

- 5 R^{3a} is selected from the group C_{1-6} alkyl, C_{1-4} cycloalkyl- C_{1-6} alkyl and benzyl, each benzyl being substituted on the aryl moiety with 0-1 substituents selected from the group C_{1-4} alkyl, Br, Cl, F, I, C_{1-4} haloalkyl, nitro, C_{1-4} alkoxy, C_{1-4} haloalkoxy, and dimethylamino;
- 10 R^4 and R^5 are independently selected at each occurrence from H, Br, Cl, F, I, -CN, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-8} cycloalkyl, C_{1-6} alkyloxy, C_{1-6} alkylthio, C_{1-6} alkylsulfinyl, C_{1-6} alkylsulfonyl, amino, C_{1-4} alkylamino,
- 15 $(C_{1-4}$ alkyl)₂ amino and phenyl, each phenyl is substituted with 0-3 groups selected from the group consisting of C_{1-7} alkyl, C_{3-8} cycloalkyl, Br, Cl, F, I, -C(O)H, C_{1-4} haloalkyl, nitro, C_{1-4} alkoxy, C_{1-4} haloalkoxy, C_{1-4} alkylthio, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, C_{1-6} alkylamino and $(C_{1-4}$ alkyl)₂ amino and wherein R^4 and R^5
- 20 non-phenyl groups may be substituted with 0-5 substituents selected from OH, halogen, -C(O)H, -OC₁₋₆-alkyl and C_{1-6} haloalkyl, C_{1-6} alkyl, C_{3-7} c-alkyl, C_{1-6} alkyl(OH)_nCO₂R wherein R is H or C_{1-6} alkyl, C_{1-6} alkyl(OH)_n,
- 25 ,wherein n is 0-3 or R^4 and R^5 may join together to form a C_{3-6} alkylene chain;


- R^6 , R^{6a} and R^7 are independently selected from:
- H, C_{1-10} alkyl, C_{3-10} cycloalkyl, C_{3-10} alkenyl, C_{3-10} alkynyl,
- 30 C_{1-10} haloalkyl, C_{2-8} alkoxyalkyl, C_{4-12} cycloalkylalkyl, C_{5-10} cycloalkenyl, C_{6-14} cycloalkenylalkyl;

- R^6 , R^{6a} and R^7 are substituted with 0-6 substituents independently selected from halogen, C_{1-4} alkyl, C_{3-8} cycloalkyl, C_{1-6} alkyloxy or C_{1-4} haloalkyl;
- 35

with the proviso that the compounds of Formula I with R^1 , R^2 , R^3 , R^4 and R^5 as specifically defined below are excluded:

- 5 (a) a compound of formula I wherein $A = CR^5$ with R^5 o-hydroxyphenyl, $B = N$, $R^3 =$ o-hydroxyphenyl, $R^1 = SMe$ and $R^2 = CN$;
- (b) a compound of formula I wherein $A = CR^5$, $R^5 = CH_3$, $B = N$,
10 $R^1 = Ph$, $R^2 = Br$ and R^3 is Ph ;
- (c) a compound of formula I wherein $A = CR^5$, $R^5 = p\text{-Cl-phenyl}$, $B = N$, $R^1 = Me$, $R^2 = H$ and $R^3 = p\text{-CF}_3\text{-phenyl}$;
- 15 (d) a compound of formula I wherein $A = CR^5$, $R^5 = \text{phenyl}$, $B = N$, $R^1 = Me$, $R^2 = H$ and $R^3 = p\text{-CF}_3\text{-phenyl}$;
- (e) a compound of formula I wherein $A = CR^5$, $R^5 = \text{ethyl}$, $B = N$, $R^1 = Me$, $R^2 = H$ and $R^3 = N\text{-methyl-piperiazin-N-yl}$;
20
- (f) a compound of formula I wherein $A = CR^5$, R^5 is $p\text{-Cl-Ph}$, $R^1 = H$, $R^2 = H$ and $R^3 = p\text{-CF}_3\text{-Ph}$;
- (g) a compound of formula I wherein $A = CR^5$, $R^5 = p\text{-Cl-Ph}$, $R^1 =$
25 CH_3 , $R^2 = H$, $R^3 = p\text{-CF}_3\text{-Ph}$;
- (h) a compound of formula I wherein $A = CR^5$, $R^5 = Ph$, $R^1 = Me$, $R^2 = H$, $R^3 = p\text{-CF}_3\text{-Ph}$;
- 30 (i) a compound of formula I wherein $A = CR^5$, $R^5 = Ph$, $R^1 = H$, $R^2 = H$, $R^3 = p\text{-CF}_3\text{-Ph}$;
- (j) a compound of formula I wherein $A = CR^5$, $R^3 = Ph$ and R^2 is H , Br , CN , CO_2Et or Cl ;
35
- (k) a compound of formula I wherein $A = CR^5$, $R^5 = CH_3$, C_2H_5 or Ph , $R^1 = H$, $R^2 = H$ and $R^3 = Ph$.

2. A compound of formula I:



5

or a stereoisomer or pharmaceutically acceptable salt thereof, wherein:

A equals N or CR⁵;

10

B equals N or CR⁴ ;

provided that both A and B cannot be N or

provided that A can not be CR⁵ and B can not be CR⁴ to

15 form a pyrazolopyrimidine; and wherein,

R¹ is independently selected from the group consisting of

H,

20 halogen,

CN,

C₁₋₆ alkyl,

C₂₋₁₀ alkenyl,

C₂₋₁₀ alkynyl,

25 C₃₋₆ cycloalkyl,

C₁₋₆ alkyloxy,

C₁₋₆ alkylS(O)_n,

-NR^{1a}R^{1b} wherein R^{1a} and R^{1b} are independently selected from

H, C₁₋₄ alkyl, C₃₋₈ cycloalkyl, -C(O)C₁₋₄alkyl,

30 C₁₋₆ alkylNR^{1a}R^{1b},

NR^{1a}COR^{1b},

-C(O)NR^{1a}R^{1b},

-O-C(O)C₁₋₄alkyl,

-XR^{1c} wherein R^{1c} is selected from H or -C₁₋₄ alkylaryl;

X is selected from O or S(O)_n,

5

wherein R¹ is substituted with 0-6 substituents selected from halogen, C₁₋₄ alkyl, C₃₋₈ cycloalkyl, C₁₋₆ alkyloxy, C₁₋₄ haloalkyl, C₁₋₄ alkylamino, C₂₋₈ dialkylamino, C₁₋₄ alkyloxy, C₁₋₄ alkylthio, C₁₋₄ alkylsulfinyl or C₁₋₄ alkylsulfonyl;

10

R² is selected from the group consisting of

OR⁷, SH, NR⁶R⁷, C(OH)R⁶R^{6a}, C(OR⁷)R⁶R^{6a}, S(O)_nR¹³, COR⁷, CO₂R⁷, CHR⁶(OR⁷)R^{6a}, OC(O)R¹³, NO, NO₂, NR⁶C(O)R⁷, N(COR⁷)₂, NR⁸CONR⁶R⁷ or NR⁶CO₂R⁷; or R² is selected from:

15

C₁₋₁₀ alkyl,

C₂₋₁₀ alkenyl,

C₂₋₁₀ alkynyl,

C₃₋₈ cycloalkyl,

20 C₃₋₆ cycloalkyl C₁₋₆ alkyl,

C₁₋₁₀ alkyloxy,

C₁₋₁₀ alkyloxyC₁₋₁₀ alkyl,

-SO₂-C₁₋₁₀alkyl

-SO₂R^{2a} wherein R^{2a} is aryl,

25 -SO₂R^{2b} wherein R^{2b} is heteroaryl,

-NR^{2c}R^{2d} wherein R^{2c} and R^{2d} are independently selected from H, C₁₋₈ alkyl, S(O)_nC₁₋₄alkyl, C(O)NR^{2c}R^{2d}, CO₂C₁₋₄alkyl, C₃₋₈ cycloalkyl, C₁₋₆ alkyloxyC₁₋₆ alkyl, -C(O)C₁₋₄alkyl or R^{2c} and R^{2d} may join to form a heterocyclic ring
30 having 0-3 heteroatoms selected from O, N or S,

-C(O)-L wherein L is selected from H, NR^{2c}R^{2d}, C₁₋₆ alkyl

O(CH₂)_mOR wherein R is C₁₋₃ alkyl, O(CH₂)_m-NR^{2c}R^{2d}, OH,

35 C(O)OC₁₋₆alkyl, or aryl or heteroaryl wherein m is 1-4; or

-OC(O)-M wherein M is selected from C₁₋₄ alkyl, C₁₋₄

haloalkyl, C₂₋₈ alkoxyalkyl, C₃₋₆cycloalkyl, C₄₋₁₂

cycloalkylalkyl, aryl, C₁₋₆ alkylaryl, heteroaryl, C₁₋₆ alkylheteroaryl;

n is 0, 1 or 2; and wherein

5

R² is substituted with 0-3 substituents independently selected from R', R'', R''' wherein R', R'' and R''' are independently selected from C₁₋₆ alkyl, C₃₋₇ cycloalkyl, hydroxyc₁₋₆ alkyl, C₁₋₆ alkyloxyC₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ alkyloxy, hydroxy, or

10

R² is substituted with 0-3 substituents independently selected from:

15 halogen,

-CN,

-S(O)_nR^{2o} wherein R^{2o} is selected from C₁₋₄ alkyl, C₁₋₄ haloalkyl, C₁₋₄ alkyloxy C₁₋₄ alkyl, C₃₋₆ cycloalkyl;

20 -COR^{2f} wherein R^{2f} is selected from H, C₁₋₄ alkyl, C₁₋₄ haloalkyl, C₁₋₄ alkyloxy C₁₋₄ alkyl, C₃₋₆ cycloalkyl, and C₃₋₆ cycloalkylC₁₋₄ alkyl;

-CO₂R^{2f},

25 -NR^{2g}COR^{2f} wherein R^{2g} is selected from H, C₁₋₆ alkyl, C₃₋₇ cycloalkyl, C₃₋₆ cycloalkyl C₁₋₆ alkyl;

-N(COR^{2f})₂,

-NR^{2g}CONR^{2f}R^{2h}, wherein R^{2h} is selected from H, C₁₋₆ alkyl, C₁₋₄ haloalkyl, C₁₋₄ alkoxy C₁₋₄ alkyl, C₃₋₆ cycloalkyl and C₃₋₆ cycloalkylC₁₋₆ alkyl;

30

-NR^{2g}CO₂R^{2o},

-CONR^{2g}R^{2h},

35 1-morpholinyl,
1-piperidinyl,
1-piperazinyl,
and

- C_{3-8} cycloalkyl wherein 0-1 carbon atoms in the C_{4-8} cycloalkyl is replaced by a group selected from -O-, $-S(O)_n-$, $-NR^{2g}-$, $-NCO_2R^{2o}$, $-NCOR^{2o}$, and $-NSO_2R^{2o}$; and wherein N_4 in
- 5 1-piperazinyl is substituted with 0-1 substituents selected from R^{2g} , CO_2R^{2o} , COR^{2o} and SO_2R^{2o} ; or
- the group R^{2i} , R^{2j} , R^{2k} , C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, Br, Cl, F, I, C_{1-4} haloalkyl, $-OR^{2g}$, $-NR^{2g}R^{2h}$, $-C_{1-6}$ alkyl- OR^{2g} , and C_{3-8} cycloalkyl which is
- 10 substituted with 0-1 R^{2i} and in which 0-1 carbons of C_{4-8} cycloalkyl is replaced by -O-, wherein
- 15 R^{2i} is selected from aryl wherein aryl includes phenyl, naphthyl, indanyl and indenyl, each R^{2i} being substituted with 0-1 OR^{2m} and 0-5 substituents independently selected from the group C_{1-6} alkyl, C_{3-6} cycloalkyl, Br, Cl, F, I, C_{1-4} haloalkyl, -CN,
- 20 nitro, -SH, $-S(O)_nR^{2n}$, $-COR^{2m}$, $-OC(O)R^{2n}$, $-NR^{2g}COR^{2m}$, $-N(COR^{2m})_2$, $-NR^{2g}CONR^{2o}R^{2p}$, $-NR^{2g}CO_2R^{2n}$, $-NR^{2o}R^{2p}$ and $-CONR^{2o}R^{2p}$;
- R^{2j} is selected from heteroaryl wherein heteroaryl
- 25 includes pyridyl, pyrimidinyl, triazinyl, furanyl, quinolinyl, isoquinolinyl, thienyl, imidazolyl, thiazolyl, indolyl, pyrrolyl, oxazolyl, benzofuranyl, benzothienyl, benzothiazolyl, benzoxazolyl, isoxazolyl, pyrazolyl, triazolyl, tetrazolyl, indazolyl, 2,3-
- 30 dihydrobenzofuranyl, 2,3-dihydrobenzothienyl, 2,3-dihydrobenzothienyl-s-oxide, 2,3-dihydro-benzothienyl-S-dioxide, indolinyl, benzoxazolin-2-onyl, benzodioxolanyl and benzodioxane, each heteroaryl being substituted on 0-4 carbon atoms with a substituent independently selected
- 35 from the group C_{1-6} alkyl, C_{3-6} cycloalkyl, Br, Cl, F, I, C_{1-4} haloalkyl, -CN, nitro, OR^{2m} , -SH, $-S(O)_nR^{2h}$, $-COR^{2m}$, $-OC(O)R^{2h}$, $-NR^{2g}COR^{2m}$, $-N(COR^{2m})_2$, $-NR^{2g}CONR^{2o}R^{2p}$, $-NR^{2g}CO_2R^{2h}$, $-NR^{2o}R^{2p}$ and $-CONR^{2o}R^{2p}$ and each heteroaryl being substituted

on any nitrogen atom with 0-1 substituents selected from the group R^{2g} , CO_2R^{2o} , COR^{2o} and SO_2R^{2o} ;

R^{2k} is heterocyclyl which is a saturated or partially saturated heteroaryl as defined for R^{2j} , each heterocyclyl being substituted on 0-4 carbon atoms with a substituent independently selected from the group C_{1-6} alkyl, C_{3-6} cycloalkyl, Br, Cl, F, I, C_{1-4} haloalkyl, -CN, nitro, $-OR^{2m}$, -SH, $-S(O)_nR^{2h}$, $-COR^{2m}$, $-OC(O)R^{2h}$, $-NR^{2g}COR^{2m}$, $-N(COR^{2m})_2$, $-NR^{2g}CONR^{2o}R^{2p}$, $NR^{2g}CO_2R^{2h}$, $-NR^{2o}R^{2p}$ and $-CONR^{2o}R^{2p}$ and each heterocyclyl being substituted on any nitrogen atom with 0-1 substituents selected from the group R^{2f} , CO_2R^{2o} , COR^{2o} and SO_2R^{2o} ;

wherein

R^{2i} is H, C_{1-4} alkyl, C_{3-6} cycloalkyl- C_{1-4} alkyl and C_{3-8} cycloalkyl;

R^{2m} is H, C_{1-6} alkyl, C_{3-6} cycloalkyl C_{1-6} alkyl, C_{1-2} alkyloxy C_{1-2} alkyl, C_{1-4} haloalkyl, $R^{2g}S(O)_n-C_{1-4}$ alkyl and $R^{2r}R^{2s}N-C_{2-4}$ alkyl;

R^{2n} is H, C_{1-6} alkyl, C_{3-10} cycloalkyl, C_{3-6} cycloalkyl- C_{1-6} alkyl, C_{1-2} alkyloxy C_{1-2} alkyl, and C_{1-4} haloalkyl;

R^{2o} and R^{2p} are independently selected at each occurrence from H, C_{1-6} alkyl, C_{3-10} cycloalkyl, C_{3-6} cycloalkyl C_{1-6} alkyl and C_{1-4} haloalkyl;

R^{2q} is selected from C_{1-6} alkyl, C_{1-4} haloalkyl, C_{1-4} alkoxy- C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkyl- C_{1-6} alkyl, aryl, aryl(C_{1-4} alkyl), heteroaryl and heteroaryl (C_{1-4} alkyl)- and benzyl, each benzyl being substituted on the aryl moiety with 0-1 substituents selected from the group C_{1-4} alkyl, Br, Cl, F, I, C_{1-4} haloalkyl, nitro, C_{1-4} alkoxy C_{1-4} haloalkoxy, and dimethylamino;

$R^{2r}R^{2v}$ taken together with the N form 1-pyrrolidinyl, 1-morpholinyl, 1-piperidinyl or 1-piperazinyl wherein N_4 in 1-piperiazinyl is substituted with 0-1 substituents selected from the group R^{2t} , CO_2R^{2q} , COR^{2q} and SO_2R^{2q} ;

5

R^{2t} is selected from H, C_{1-4} alkyl, C_{1-4} haloalkyl, C_{1-4} alkoxy- C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkyl - C_{1-6} alkyl, aryl, aryl (C_{1-4} alkyl)-, heteroaryl and heteroaryl (C_{1-4} alkyl);

10

R^3 is selected from an aryl or heteroaryl group attached through an unsaturated carbon atom;

aryl is selected from phenyl, naphthyl, indanyl and indenyl, each aryl being substituted with 0-5 substituents independently selected at each occurrence from C_{1-6} alkyl, C_{3-6} cycloalkyl, methylenedioxy, C_{1-4} alkyloxy- C_{1-4} alkyloxy, $-OR^{2m}$, Br, Cl, F, I, C_{1-4} haloalkyl, -CN, $-NO_2$, -SH, $-S(O)_nR^{2n}$, $-COR^{2m}$, $-CO_2R^{2m}$, $-OC(O)R^{2n}$, -
 20 $NR^{2g}COR^{2m}$, $-N(COR^{2m})_2$, $-NR^{2g}CONR^{2o}R^{2p}$, $-NR^{2g}CO_2R^{2h}$, $-NR^{2o}R^{2p}$ and $CONR^{2o}R^{2p}$;

heteroaryl is selected from the group pyridyl, pyrimidyl, triazinyl, furanyl, quinolinyl, isoquinolinyl, thienyl, imidazolyl, thiazolyl, indolyl, pyrrolyl, oxazolyl, benzofuranyl, benzothienyl, benzothiazolyl, benzoxazolyl, isoxazolyl, triazolyl, tetrazolyl, indazolyl, 2,3-dihydrobenzo-furanyl, 2,3-dihydrobenzothienyl, 2,3-dihydro-benzothienyl-S-oxide, 2,3-dihydrobenzothienyl-s-dioxide, indolinyl, benzoxazolin-2-on-yl, benzodioxolanyl and benzodioxane, each heteroaryl being substituted at 0-4 carbon atoms with a substituent independently selected at each occurrence from the group C_{1-6} alkyl, C_{3-6} cycloalkyl, Br, F, I, C_{1-4} haloalkyl, -CN, $NR^{2g}R^{2h}$, nitro, -
 35 OR^{2n} , -SH, $-S(O)_nR^{2n}$, COR^{2m} , $-CO_2R^{2m}$, $-OC(O)R^{2n}$, $-NR^{2g}COR^{2m}$, -
 $N(COR^{2m})_2$, $-NR^{2g}CONR^{2o}R^{2p}$ and each heteroaryl being substituted at any nitrogen atom with 0-1 substituents selected from the group R^{2g} , CO_2R^{3a} , COR^{3a} and SO_2R^{3a} wherein,



R^{3a} is selected from the group C_{1-6} alkyl, C_{1-4} cycloalkyl- C_{1-6} alkyl and benzyl, each benzyl being substituted on the aryl moiety with 0-1 substituents selected from the group
5 C_{1-4} alkyl, Br, Cl, F, I, C_{1-4} haloalkyl, nitro, C_{1-4} alkoxy, C_{1-4} haloalkoxy, and dimethylamino;

R^4 and R^5 are independently selected at each occurrence from H, Br, Cl, F, I, -CN, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-8} cycloalkyl, C_{1-6} alkyloxy, C_{1-6} alkylthio, C_{1-6} alkylsulfinyl, C_{1-6} alkylsulfonyl, amino, C_{1-4} alkylamino, $(C_{1-4}$ alkyl)₂ amino and phenyl, each phenyl is substituted with 0-3 groups selected from the group consisting of C_{1-7} alkyl, C_{3-8} cycloalkyl, Br, Cl, F, I, -C(O)H, C_{1-4} haloalkyl, nitro, C_{1-4} alkoxy, C_{1-4} haloalkoxy, C_{1-4} alkylthio, C_{1-4} alkylsulfinyl, C_{1-4} alkylsulfonyl, C_{1-6} alkylamino and $(C_{1-4}$ alkyl)₂ amino and wherein R^4 and R^5 non-phenyl groups may be substituted with 0-5 substituents selected from OH, halogen, -C(O)H, -OC₁₋₆- alkyl and C_{1-6} haloalkyl, C_{1-6} alkyl, C_{3-7} c-alkyl, C_{1-6} alkyl(OH)_nCO₂R wherein R is H or C_{1-6} alkyl, C_{1-6} alkyl(OH)_n, wherein n is 0-3 or R^4 and R^5 may join together to form a C_{3-6} alkylene chain;

25 R^6 , R^{6a} and R^7 are independently selected from:
H, C_{1-10} alkyl, C_{3-10} cycloalkyl, C_{3-10} alkenyl, C_{3-10} alkynyl, C_{1-10} haloalkyl, C_{2-8} alkoxyalkyl, C_{4-12} cycloalkylalkyl, C_{5-10} cycloalkenyl, C_{6-14} cycloalkenylalkyl;

30 R^6 , R^{6a} and R^7 are substituted with 0-6 substituents independently selected from halogen, C_{1-4} alkyl, C_{3-8} cycloalkyl, C_{1-6} alkyloxy, C_{1-4} haloalkyl.

35 3. A compound of formula (Ia) or (Ib)



wherein R^1 - R^5 are as defined in Claims 1 or 2.

5

4. The compound according to Claim 1, 2 or 3 wherein

R^1 is selected from C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{1-6} alkoxy, C_{1-6} alkylthio, $-XR^{1c}$ wherein R^1 is substituted with
 10 0-6 substituents selected from halogen, C_{1-4} alkyl or C_{1-4} haloalkyl;


R^2 is selected from C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{3-8} cycloalkyl, C_{3-6} cycloalkyl C_{1-6} alkyl, and -
 15 $NR^{2c}R^{2d}$ wherein R^2 is unsubstituted or substituted with 1-3 substituents independently selected from the group R^{2i} , R^{2j} , R^{2k} , C_{1-6} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, Br, Cl, F, I, C_{1-4} haloalkyl, $-OR^{2g}$, $-NR^{2g}R^{2h}$, $-C_{1-6}$ alkyl- OR^{2g} , and C_{3-8} cycloalkyl which is substituted with 0-1 R^{2i} and in which
 20 0-1 carbons of C_{4-8} cycloalkyl is replaced by -O-.

5. The compound according to Claims 1, 2, 3 or 4 wherein R^3 is selected from an aryl group selected from phenyl or substituted versions thereof or a heteroaryl
 25 group selected from pyridyl or substituted versions thereof.

6. The compounds according to Claims 1, 2, 3, 4 or 5 wherein R^3 is substituted with 0-4 substituents
 30 independently selected from halogen, C_{1-4} alkoxy, C_{1-6} alkyl or $NR'R''$ wherein R' and R'' are independently selected from H or C_{1-6} alkyl.

7. A compound of formula (Ia)

5



(Ia)

10 or a pharmaceutically acceptable salt thereof, wherein

15 R^1 is independently selected at each occurrence from H, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, halo, CN, C₁-C₄ haloalkyl, C₁-C₁₂ hydroxyalkyl, C₂-C₁₂ alkoxyalkyl, C₂-C₁₀ cyanoalkyl, C₃-C₆ cycloalkyl, C₄-C₁₀ cycloalkylalkyl, NR⁹R¹⁰, C₁-C₄ alkyl-NR⁹R¹⁰, NR⁹COR¹⁰, OR¹¹, SH or S(O)_nR¹²;

R^2 is selected from:

20 -H, OR⁷, SH, S(O)_nR¹³, COR⁷, CO₂R⁷, CHR⁶(OR⁷)R^{6a}, OC(O)R¹³, CH(OH)R⁶, C(OH)R⁶R^{6a}, C(OR⁷)R⁶R^{6a}, NO, NO₂, NR⁶COR⁷, N(COR⁷)₂, NR⁸CONR⁶R⁷, NR⁶CO₂R⁷, NR⁶R⁷, NR⁶S(O)₂R⁷, N(S(O)₂R⁷)₂, N(OR⁷)R⁶, CONR⁶R⁷;

25 or

-C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₃-C₈ cycloalkyl, C₅-C₈ cycloalkenyl, C₄-C₁₂ cycloalkylalkyl or C₆-C₁₀ cycloalkenylalkyl, each optionally substituted with 1 to 3 substituents independently selected at each occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo,

30

C₁-C₄ haloalkyl, cyano, OR¹⁵, SH,
 S(O)_nR¹³, COR¹⁵, CO₂R¹⁵, OC(O)R¹³, NR⁸COR¹⁵,
 N(COR¹⁵)₂, NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹³, NR¹⁶R¹⁵,
 CONR¹⁶R¹⁵;

5

R³ is selected from phenyl, naphthyl, pyridyl,
 pyrimidinyl, triazinyl, furanyl, thienyl,
 benzothienyl, benzofuranyl, 2,3-
 dihydrobenzofuranyl, 2,3-dihydrobenzothienyl,
 10 indanyl, 1,2-benzopyranyl, 3,4-dihydro-1,2-
 benzopyranyl, tetralinyl, each R³ optionally
 substituted with 1 to 5 substituents and each Ar
 is attached via an unsaturated carbon atom wherein
 the substituents are independently selected at
 15 each occurrence from: C₁-C₁₀ alkyl, C₂-
 C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₃-C₆ cycloalkyl, C₄-
 C₁₂ cycloalkylalkyl, NO₂, halo, CN, C₁-
 C₄ haloalkyl, NR⁶R⁷, NR⁸COR⁷, NR⁸CO₂R⁷, COR⁷, OR⁷,
 CONR⁶R⁷, CO(NOR⁹)R⁷, CO₂R⁷, or S(O)_nR⁷, where each
 20 such C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl,
 C₃-C₆ cycloalkyl and C₄-C₁₂ cycloalkylalkyl are
 optionally substituted with 1 to 3 substituents
 independently selected at each occurrence from C₁-
 C₄ alkyl, NO₂, halo, CN, NR⁶R⁷, NR⁶COR⁷, NR⁷CO₂R⁷,
 25 COR⁷ OR⁷, CONR⁶R⁷, CO₂R⁷, CO(NOR⁹)R⁷, or S(O)_nR⁷;

R⁵ is selected from H, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-
 C₄ alkynyl, C₃-C₆ cycloalkyl, C₄-C₁₀
 cycloalkylalkyl, each optionally substituted with
 1 to 3 substituents independently selected at
 30 each occurrence from C₁-C₆ alkyl, C₃-
 C₆ cycloalkyl; halo, C₁-C₄ haloalkyl, cyano, OR¹⁵,
 SH, S(O)_nR¹³, COR¹⁵, CO₂R¹⁵, OC(O)R¹³, NR⁸COR¹⁵,
 N(COR¹⁵)₂, NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹³, NR¹⁶R¹⁵,
 35 CONR¹⁶R¹⁵, aryl, heteroaryl and heterocyclyl;
 or

halo, CN, $-NR^6R^7$, NR^9COR^{10} , $-NR^6S(O)_nR^7$,
 $S(O)_nNR^6R^7$, C₁-C₄ haloalkyl, $-OR^7$, SH or -
 $S(O)_nR^{12}$;

5

R^6 , R^{6a} and R^7 are independently selected at each
occurrence from:

-H,

10 -C₁-C₁₀ alkyl, C₃-C₁₀ alkenyl, C₃-C₁₀ alkynyl,
C₁-C₁₀ haloalkyl with 1-10 halogens, C₂-C₈
alkoxyalkyl, C₃-C₆ cycloalkyl, C₄-
C₁₂ cycloalkylalkyl, C₅-C₁₀ cycloalkenyl,
or C₆-C₁₄ cycloalkenylalkyl, each optionally
15 substituted with 1 to 3 substituents
independently selected at each occurrence from
C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-
C₄ haloalkyl, cyano, OR^{15} , SH, $S(O)_nR^{13}$, COR^{15} ,
 CO_2R^{15} , $OC(O)R^{13}$, NR^8COR^{15} , $N(COR^{15})_2$,
20 $NR^8CONR^{16}R^{15}$, $NR^8CO_2R^{13}$, $NR^{16}R^{15}$, $CONR^{16}R^{15}$,
aryl, heteroaryl or heterocyclyl,
-aryl, aryl(C₁-C₄ alkyl), heteroaryl,
heteroaryl(C₁-C₄ alkyl), heterocyclyl or
heterocyclyl(C₁-C₄ alkyl);

25

alternatively, NR^6R^7 and $NR^{6a}R^{7a}$ are independently
piperidine, pyrrolidine, piperazine, N-
methylpiperazine, morpholine or thiomorpholine, each
optionally substituted with 1-3 C₁-C₄ alkyl groups;

30

R^8 is independently selected at each occurrence from H
or C₁-C₄ alkyl;

35 R^9 and R^{10} are independently selected at each
occurrence from H, C₁-C₄ alkyl, or C₃-C₆
cycloalkyl;

R¹¹ is selected from H, C₁-C₄ alkyl, C₁-C₄ haloalkyl, or C₃-C₆ cycloalkyl;

R¹² is C₁-C₄ alkyl or C₁-C₄ haloalkyl;

5

R¹³ is selected from C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₂-C₈ alkoxyalkyl, C₃-C₆ cycloalkyl, C₄-C₁₂ cycloalkylalkyl, aryl, aryl(C₁-C₄ alkyl)-, heteroaryl or heteroaryl(C₁-C₄ alkyl)-;

10

R¹⁵ and R¹⁶ are independently selected at each occurrence from H, C₁-C₆ alkyl, C₃-C₁₀ cycloalkyl, C₄-C₁₆ cycloalkylalkyl, except that for S(O)_nR¹⁵, R¹⁵ cannot be H;

15

aryl is phenyl or naphthyl, each optionally substituted with 1 to 5 substituents independently selected at each occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹⁵, COR¹⁵, CO₂R¹⁵, OC(O)R¹⁵, NR⁸COR¹⁵, N(COR¹⁵)₂, NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹⁵, NR¹⁶R¹⁵, and CONR¹⁶R¹⁵;

20

heteroaryl is pyridyl, pyrimidinyl, triazinyl, furanyl, pyranlyl, quinolinyl, isoquinolinyl, thienyl, imidazolyl, thiazolyl, indolyl, pyrrolyl, oxazolyl, benzofuranyl, benzothienyl, benzothiazolyl, isoxazolyl, pyrazolyl, 2,3-dihydrobenzothienyl or 2,3-dihydrobenzofuranyl, each being optionally substituted with 1 to 5 substituents independently selected at each occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹⁵, -COR¹⁵, CO₂R¹⁵, OC(O)R¹⁵, NR⁸COR¹⁵, N(COR¹⁵)₂, NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹⁵, NR¹⁶R¹⁵, and CONR¹⁶R¹⁵;

25

30

35

heterocyclyl is saturated or partially saturated heteroaryl, optionally substituted with 1 to 5


substituents independently selected at each occurrence from C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halo, C₁-C₄ haloalkyl, cyano, OR¹⁵, SH, S(O)_nR¹⁵, COR¹⁵, CO₂R¹⁵, OC(O)R¹⁵, NR⁸COR¹⁵, N(COR¹⁵)₂,
 5 NR⁸CONR¹⁶R¹⁵, NR⁸CO₂R¹⁵, NR¹⁵R¹⁶, and CONR¹⁶R¹⁵;

n is independently at each occurrence 0, 1 or 2.

- 10 8. The compound according to Claims 1-7 wherein R² is selected from 3-pentyl, NEt₂, butyl, NHCH(CH₂OMe)₂, NHCH(CH₂OEt)₂, NHCH(Et)CH₂OMe, NH-3-heptyl, NH-3-pentyl, NH-2-butyl, NH-3-hexyl, NHCH(CH₂Ph)CH₂OMe, NHCH(Et)CH₂CH₂OMe, NH-cyclobutyl, NH-cyclopentyl, NEtPr,
 15 NEtBu, NMePr, NMePh, Npr₂, NPr(CH₂-c-C₃H₅), N(CH₂CH₂OMe)₂, morpholino, N(CH₂Ph)CH₂CH₂OMe, N(Me)CH₂CH₂OMe, N(Et)CH₂CH₂OMe, N(CH₂-c-C₃H₅)CH₂CH₂OMe, N(CH₂-c-C₃H₅)Pr, N(CH₂-c-C₃H₅)Et, OEt, OCH(Et)CH₂OMe, OCH(Et)CH₂CH₂OMe, OCH(Me)CH₂CH₂OMe, O-3-pentyl, O-2-pentyl, S-3-pentyl, S-2-pentyl, SEt, S(O)Et, SO₂Et, S-3-pentyl, S(O)-3-pentyl, SO₂-3-pentyl, S-2-pentyl, S(O)-2-pentyl, SO₂-2-pentyl, CH(CO₂Et)₂, C(Et)(CO₂Et)₂, CH(Et)CH₂OH, CH(Et)CH₂OMe, CH(Et)CH₂CH₂OMe, CONMe₂, COCH₃, COEt, COPr, CO-2-pentyl, CO-3-pentyl, CH(OH)CH₃,
 25 C(OH)Me₂, C(OH)Ph-3-pyridyl, CH(OMe)CH₃, CH(OMe)Et, CH(OMe)Pr, CH(OEt)CH₃, CH(OPr)CH₃, 2-pentyl, 2-butyl, cyclobutyl, cyclopentyl, CH(Me)cyclobutyl, CH(OMe)cyclobutyl, CH(OH)cyclobutyl, CH(Me)cyclopropyl, CH(OMe)cyclopropyl, CH(OH)cyclopropyl, CH(Et)cyclobutyl, CH(Et)cyclopropyl, CH(OMe)cyclobutyl, CH(OMe)cyclopropyl, CH(OEt)cyclobutyl, CH(OEt)cyclopropyl, CH(Me)CH₂-cyclobutyl, CH(OMe)CH₂-cyclobutyl, CH(OH)CH₂-cyclobutyl, CH(Me)CH₂-cyclopropyl, CH(OMe)CH₂-cyclopropyl, CH(OH)CH₂-cyclopropyl, CH(Et)CH₂-cyclobutyl, CH(Et)CH₂-cyclopropyl,
 30 CH(OMe)CH₂-cyclobutyl, CH(OMe)CH₂-cyclopropyl, CH(OEt)CH₂-cyclobutyl, CH(OEt)CH₂-cyclopropyl,

- CH(CH₂OMe)cyclobutyl, CH(CH₂OMe)cyclopropyl,
 CH(CH₂OEt)cyclobutyl, CH(CH₂OEt)cyclopropyl,
 CH(cyclobutyl)₂, CH(cyclopropyl)₂, CH(Et)CH₂CONMe₂,
 CH(Et)CH₂CH₂NMe₂, CH(CH₂OMe)Me, CH(CH₂OMe)Et,
 5 CH(CH₂OMe)Pr, CH(CH₂OEt)Me, CH(CH₂OEt)Et, CH(CH₂OEt)Pr,
 CH(CH₂C≡CMe)Et, CH(CH₂C≡CMe)Et.


9. A compound of formula Ib



10

having R¹-R⁴ as defined in Claims 1-8.

10. A compound of formula II



- 15 or a pharmaceutically acceptable salt or isomer thereof
 wherein R¹-R⁴ are as defined in any of claims 1-8.

11. Use of a compound according to Claims 1-10 in
 therapy.

20

12. Use of a compound according to Claims 1-10 to
 antagonize a CRF-1 receptor in mammals including humans
 wherein binding to the receptor causes and ultimately
 results in the treatment of affective disorder, anxiety,
 25 depression, headache, irritable bowel syndrome, post-
 traumatic stress disorder, supranuclear palsy, immune
 suppression, Alzheimer's disease, gastrointestinal

diseases, anorexia nervosa or other feeding disorder,
drug addiction, drug or alcohol withdrawal symptoms,
inflammatory diseases, cardiovascular or heart-related
diseases, fertility problems, human immunodeficiency
5 virus infections, hemorrhagic stress, obesity,
infertility, head and spinal cord traumas, epilepsy,
stroke, ulcers, amyotrophic lateral sclerosis,
hypoglycemia or a disorder the treatment of which can be
effected or facilitated by antagonizing CRF, including
10 but not limited to disorders induced or facilitated by
CRF, in mammals comprising administering to the mammal a
therapeutically effective amount of a compound according
to Claims 1-10 with the proviso that, in the case of
compounds of Claim 1, the provisos are not present.

15

13. A pharmaceutical composition comprising a compound
according to Claims 1-10 and a pharmaceutically
acceptable carrier.

20

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
12 October 2000 (12.10.2000)

PCT

(10) International Publication Number
WO 00/59907 A3

- (51) International Patent Classification⁷: C07D 487/04, A61K 31/53, A61P 25/00, 9/00 // (C07D 487/04, 251:00, 231:00) (C07D 487/04, 253:00, 231:00)
- (74) Agent: RUBIN, Kenneth, B.; Du Pont Pharmaceuticals Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).
- (21) International Application Number: PCT/US00/09109
- (81) Designated States (*national*): AU, BR, CA, CN, CZ, EE, HU, IL, IN, JP, KR, LT, LV, MX, NO, NZ, PL, RO, SG, SI, SK, TR, UA, VN, ZA.
- (22) International Filing Date: 6 April 2000 (06.04.2000)
- (25) Filing Language: English
- (84) Designated States (*regional*): Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
- (26) Publication Language: English
- (30) Priority Data: 60/128,008 6 April 1999 (06.04.1999) US
- Published:
— *With international search report.*
- (71) Applicant: DU PONT PHARMACEUTICALS COMPANY [US/US]; Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19807 (US).
- (88) Date of publication of the international search report:
4 January 2001
- (72) Inventors: GILLIGAN, Paul, J.; 2629 Pennington Drive, Wilmington, DE 19810 (US). WILDE, Richard, G.; 205 Roseman Court, Newark, DE 19711 (US).
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: PYRAZOLOTRIAZINES AS CRF ANTAGONISTS



(57) Abstract: The present invention relates to pyrazolotriazines according to formula (I), and stereoisomers, isomers and salts thereof wherein R¹-R⁵ are selected from certain alkyl, aryl and heteroaryl species as defined in the specification wherein all of the compounds are useful as CRF antagonists and are thus useful in the treatment of neurological disorders as well as a multitude of other CRF associated diseases or conditions.

WO 00/59907 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/09109

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D487/04 A61K31/53 A61P25/00 A61P9/00
 //(C07D487/04, 251:00, 231:00), (C07D487/04, 253:00, 231:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|---|-----------------------|
| X | <p>DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; DEEB, ALI ET AL: "Preparation of naphtho'2,1-e!pyrazolo'5,1-c!'1,2,4!triazine, dipyrazolo'5,1-c: 3',4'-e!'1,2,4!triazines, and pyrazolo'1,5- c!'1,2,4!triazine derivatives" retrieved from STN Database accession no. 114:122289 XP002148012 RN 132584-89-5 & COLLECT. CZECH. CHEM. COMMUN. (1990), 55(11), 2790-4 ,</p> <p style="text-align: center;">--- -/--</p> | 1 |

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

25 September 2000

Date of mailing of the international search report

13/10/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Alfaro Faus, I

INTERNATIONAL SEARCH REPORT

National Application No

PCT/US 00/09109

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|----------|---|-----------------------|
| X | <p>DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; RAMIZ, MAHMOUD M. M. ET AL: "Studies on amino-azoles: synthesis of 3-methylaminopyrazole derivatives" retrieved from STN Database accession no. 112:35748 XP002148013 RN 124612-21-1 & ARCH. PHARM. (WEINHEIM, GER.) (1989), 322(9), 557-60 ,</p> | 1 |
| X | <p>--- DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; KANDEEL, EZZAT M. ET AL: "Reactions with heterocyclic amidines, XI. Syntheses of new 2-aminopyrazolo'1,5-a!pyrimidines and 2-amino'1,5-c!-as-triazines" retrieved from STN Database accession no. 99:122411 XP002148014 RN 87031-23-0 & ARCH. PHARM. (WEINHEIM, GER.) (1983), 316(8), 713-18 ,</p> | 1 |
| X | <p>--- DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; EGE, GUENTER ET AL: "Reactions with diazoazoles. Part IV. '7 + 2!- and '11 + 2!-Cyclocondensation reactions of diazoazoles with acyltriphenylphosphonium methylides to azolo'5,1-c!'1,2,4!triazines" retrieved from STN Database accession no. 95:169140 XP002148015 RN 79441-90-0 & J. HETEROCYCL. CHEM. (1981), 18(4), 675-7 ,</p> <p>--- -/--</p> | 1 |

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/09109

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|----------|---|-----------------------|
| X | <p>DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; GRAY, ELIZABETH JANE ET AL: "Triazines and related products. Part 21. Cyclization of 3-amino-5-hydrazinopyrazole and 3-amino-5-hydrazino-1,2,4-triazole to azolo'5,1-c!'1,2,4!triazines" retrieved from STN Database accession no. 89:215364 XP002148016 abstract & J. CHEM. SOC., PERKIN TRANS. 1 (1978), (8), 885-8 ,</p> | 1 |
| X | <p style="text-align: center;">---</p> <p>STROHMEYER, TIMOTHY W. ET AL: "New synthesis of 2,4-dialkyl(or diaryl)pyrazolo'1,5-a!-1,3,5- triazines" JOURNAL OF HETEROCYCLIC CHEMISTRY., vol. 22, no. 7, 1985, pages 7-10, XP002148011 HETEROCORPORATION. PROVO., US ISSN: 0022-152X table II, compounds 5f-5i</p> | 1 |
| A | <p style="text-align: center;">---</p> <p>WO 98 03510 A (DU PONT MERCK PHARMA) 29 January 1998 (1998-01-29) cited in the application claim 1</p> | 12 |
| P,A | <p style="text-align: center;">---</p> <p>WO 99 38868 A (DU PONT PHARM CO) 5 August 1999 (1999-08-05) claims 1,7</p> <p style="text-align: center;">-----</p> | 12 |

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/09109

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| WO 9803510 A | 29-01-1998 | AU 3894297 A | 10-02-1998 |
| | | BR 9710544 A | 17-08-1999 |
| | | CA 2259583 A | 29-01-1998 |
| | | CN 1225637 A | 11-08-1999 |
| | | CZ 9900184 A | 17-11-1999 |
| | | EP 0915880 A | 19-05-1999 |
| | | HR 970413 A | 31-10-1998 |
| | | LT 99008 A,B | 27-03-2000 |
| | | LV 12292 A | 20-06-1999 |
| | | LV 12292 B | 20-11-1999 |
| | | NO 990264 A | 10-03-1999 |
| | | PL 331523 A | 19-07-1999 |
| | | SI 9720045 A | 31-10-1999 |
| | | ZA 9706603 A | 25-01-1999 |
| WO 9938868 A | 05-08-1999 | US 6060478 A | 09-05-2000 |
| | | AU 2478799 A | 16-08-1999 |